

Weak amenability of $(2N)$ -th dual of a Banach algebra

Mina Ettefagh^{a,*}, and Sima Houdfar ^a

^a*Department of Mathematics, Tabriz Branch, Islamic Azad University, Tabriz, Iran*

Abstract. In this paper by using some conditions, we show that the weak amenability of $(2n)$ -th dual of a Banach algebra A for some $n \geq 1$ implies the weak amenability of A .

Keywords: Banach algebra, Arens products, Arens regularity, Derivation, weak amenability.

1. Introduction and Preliminaries

Let X be a normed space and X' be the topological dual space of X ; the value of $f \in X'$ at $x \in X$ is denoted by $\langle f, x \rangle$. We set $(X')' = X''$ and so on, and we regard X as a subspace of X'' by natural mapping $\iota : X \rightarrow X'' (x \mapsto \hat{x})$ where $\langle \hat{x}, f \rangle = \langle f, x \rangle (f \in X')$. We denote the n -th dual of X by $X^{(n)}$. The weak topology on X is denoted by $w = \sigma(X, X')$ and weak*-topology on X' is denoted by $w^* = \sigma(X', X)$.

Now let X, Y and Z be normed spaces and let $f : X \times Y \rightarrow Z$ be a continuous bilinear map. Arens in [1] offers two extensions f^{***} and f^{t***t} of f from $X'' \times Y''$ to Z'' as following:

$$\begin{aligned}
 (1) \quad & \begin{cases} f^* : Z' \times X \longrightarrow Y' \\ \langle f^*(z', x), y \rangle = \langle z', f(x, y) \rangle \quad (x \in X, y \in Y, z' \in Z'). \end{cases} \\
 (2) \quad & \begin{cases} f^{**} : Y'' \times Z' \longrightarrow X' \\ \langle f^{**}(y'', z'), x \rangle = \langle y'', f^*(z', x) \rangle \quad (x \in X, z' \in Z', y'' \in Y''). \end{cases} \\
 (3) \quad & \begin{cases} f^{***} : X'' \times Y'' \longrightarrow Z'' \\ \langle f^{***}(x'', y''), z' \rangle = \langle x'', f^{**}(y'', z') \rangle \quad (z' \in Z', x'' \in X'', y'' \in Y''). \end{cases}
 \end{aligned}$$

*Corresponding author. Email: minaettefagh@gmail.com

The mapping f^{***} is the unique extension of f such that $x'' \mapsto f^{***}(x'', y'')$ from X'' into Z'' is $w^* - w^*$ -continuous for every $y'' \in Y''$.

Let now $f^t : Y \times X \rightarrow Z$ be the transpose of f defined by $f^t(y, x) = f(x, y)$ for $x \in X$ and $y \in Y$. We can extend f^t as above to f^{t***} and then we have the mapping $f^{t***} : X'' \times Y'' \rightarrow Z''$. If $f^{***} = f^{t***}$ then f is called *Arens regular*. The mapping $y'' \mapsto f^{t***}(x'', y'')$ from Y'' into Z'' is $w^* - w^*$ -continuous for every $x'' \in X''$. Arens regularity of f is equivalent to the following

$$\lim_i \lim_j \langle z', f(x_i, y_j) \rangle = \lim_j \lim_i \langle z', f(x_i, y_j) \rangle,$$

whenever both limits exist for all bounded nets (x_i) and (y_j) in X and Y , respectively and for every $z' \in Z'$.

Throughout this paper A is a Banach algebra. This algebra is called *Arens regular* if its multiplication as a bilinear map $\pi : A \times A \rightarrow A (\pi(a, b) = ab)$ is Arens regular. We shall frequently use Goldstine's theorem: for each $a'' \in A''$, there is a net (a_i) in A such that $a'' = w^* - \lim_i \widehat{a_i}$. Now let $a'' = w^* - \lim_i \widehat{a_i}$ and $b'' = w^* - \lim_j \widehat{b_j}$ be elements of A'' . The first and second *Arens products* on A'' are denoted by symbols \square and \diamond respectively and defined by

$$a'' \square b'' = \pi^{***}(a'', b'') \quad , \quad a'' \diamond b'' = \pi^{t***}(a'', b'').$$

It is easy to show that

$$a'' \square b'' = w^* - \lim_i w^* - \lim_j \widehat{a_i b_j} \quad , \quad a'' \diamond b'' = w^* - \lim_j w^* - \lim_i \widehat{a_i b_j}.$$

On the other hand we can define above Arens products in stages as following. Let $a, b \in A, f \in A'$ and $F, G \in A''$.

- (1) Define $f.a$ in A' by $\langle f.a, b \rangle = \langle f, ab \rangle$,
and $a.f$ in A' by $\langle a.f, b \rangle = \langle f, ba \rangle$.
- (2) Define $F.f$ in A' by $\langle F.f, a \rangle = \langle F, f.a \rangle$,
and $f.F$ in A' by $\langle f.F, a \rangle = \langle F, a.f \rangle$.
- (3) Define $F \square G$ in A'' by $\langle F \square G, f \rangle = \langle F, G.f \rangle$,
and $F \diamond G$ in A'' by $\langle F \diamond G, f \rangle = \langle G, f.F \rangle$.

Then (A'', \square) and (A'', \diamond) are Banach algebras, see [1, 5] for further details.

Now let E be a Banach A -bimodule, then E' is a Banach A -bimodule under actions

$$\langle a.f, x \rangle = \langle f, xa \rangle, \langle f.a, x \rangle = \langle f, ax \rangle \quad (a \in A, x \in E, f \in E'), \tag{1}$$

and E'' is a Banach A'' -bimodule under actions

$$F.\Lambda = w^* - \lim_i w^* - \lim_j \widehat{a_i x_j} \quad , \quad \Lambda.F = w^* - \lim_j w^* - \lim_i \widehat{x_j a_i} \tag{2}$$

where $F = w^* - \lim_i \hat{a}_i$ and $\Lambda = w^* - \lim_j \hat{x}_j$ such that $(a_i) \subset A$ and $(x_j) \subset E$ are bounded nets.

For a Banach A -bimodule E , the continuous linear map $D : A \rightarrow E$ is called *derivation* if $D(ab) = a.D(b) + D(a).b, (a, b \in A)$. For $x \in E$ the derivation $\delta_x : A \rightarrow E$ by $\delta_x(a) = a.x - x.a$ is called *inner derivation*. The Banach algebra A is called *amenable* if every derivation $D : A \rightarrow E'$ is inner, for each Banach A -bimodule E , [7]. If every derivation $D : A \rightarrow A'$ is inner, A is called *weakly amenable*, see also [2, 4] for details.

THEOREM 1.1 *Let A be a Banach algebra and E be a Banach A -bimodule and $D : A \rightarrow E$ is a continuous derivation, then $D'' : A'' \rightarrow E''$ is a continuous derivation[5, Theorem 2.7.17].*

Remark 1 A'' -bimodule structures on E'' in above theorem are as in formula (2).

In [8] it was shown that if A is complete Arens regular and every derivation $D : A \rightarrow A'$ be weakly compact, then weak amenability of $A^{(2n)}$ for some $(n \geq 1)$ implies weak amenability of A . In this paper we always use the first Arens product \square on Banach algebra $A^{(2n)}(n \geq 1)$. In section 2 we shall frequently use formulas (1) and (2) and we investigate following actions

- ▷ two A'' -module actions on $A^{(3)} = (A')''$ and $A^{(3)} = (A'')'$,
- ▷ two $A^{(4)}$ -module actions on $A^{(5)} = ((A')'')''$ and $A^{(5)} = ((A'')'')$,
- ▷ two $A^{(6)}$ -module actions on $A^{(7)} = (((A')'')'')$ and on $A^{(7)} = (((A'')'')'')$,

and we will extend our results to two different $A^{(2n)}$ -module actions on $A^{(2n+1)}$ by induction. In each case we find conditions to make these two different actions equal. In a similar work in [6] two different A'' -module actions on $A^{(3)} = (A')''$ and $A^{(3)} = (A'')'$ have been studied. Finally in section 3 we investigate the innerness of second, fourth... and $(2n)$ -th dual of a derivation $D : A \rightarrow A'$. By using some conditions we will show that weak amenability of $A^{(2n)}$ for some $(n \geq 1)$ implies weak amenability of A .

2. $A^{(2n)}$ -module actions on $A^{(2n+1)}$

We shall frequently use formulas (1) and (2) to construct two different $A^{(2n)}$ -module actions on $A^{(2n+1)} = (((A')'') \dots)''$ and $A^{(2n+1)} = (((A'') \dots)')'$.

Remark 1 There are many other $A^{(2n)}$ -module actions on $A^{(2n+1)}$ that we don't need to mention.

First for $n=1$ we consider two A'' -module actions on $A^{(3)} = (A')''$ and $A^{(3)} = (A'')'$. Let $a^{(3)} = w^* - \lim_{\alpha} \widehat{a}'_{\alpha} \in A^{(3)}$ and $a'' = w^* - \lim_{\beta} \widehat{a}_{\beta}, b'' = w^* - \lim_i \widehat{b}_i$ in which (a'_{α}) and $(a_{\beta}), (b_i)$ are bounded nets in A' and A respectively. For left

A'' -module action on $A^{(3)} = (A')''$ as second dual of A' we can write

$$\begin{aligned} \langle a'' \cdot a^{(3)}, b'' \rangle &= \lim_{\beta} \lim_{\alpha} \langle b'', a_{\beta} \cdot a'_{\alpha} \rangle && \text{(by formula (2))} \\ &= \lim_{\beta} \lim_{\alpha} \lim_i \langle a'_{\alpha}, b_i \cdot a_{\beta} \rangle, && (3) \end{aligned}$$

and for left A'' -module action on $A^{(3)} = (A'')'$ as dual of A'' we can write

$$\begin{aligned} \langle a'' \cdot a^{(3)}, b'' \rangle &= \langle a^{(3)}, b'' \square a'' \rangle && \text{(by formula (1))} \\ &= \lim_{\alpha} \langle b'' \square a'', a'_{\alpha} \rangle && (4) \\ &= \lim_{\alpha} \lim_i \lim_{\beta} \langle a'_{\alpha}, b_i \cdot a_{\beta} \rangle. \end{aligned}$$

This shows that two left A'' -module actions on $A^{(3)} = (A'')'$ and $A^{(3)} = (A')''$ are not equal. Similarly for right A'' -module action on $A^{(3)} = (A')''$ we have

$$\begin{aligned} \langle a^{(3)} \cdot a'', b'' \rangle &= \lim_{\alpha} \lim_{\beta} \langle b'', a'_{\alpha} \cdot a_{\beta} \rangle && \text{(by formula (2))} \\ &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a'_{\alpha}, a_{\beta} \cdot b_i \rangle, && (5) \end{aligned}$$

and for right A'' -module action coincide on $A^{(3)} = (A'')'$

$$\begin{aligned} \langle a^{(3)} \cdot a'', b'' \rangle &= \langle a^{(3)}, a'' \square b'' \rangle && \text{(by formula (1))} \\ &= \lim_{\alpha} \langle a'' \square b'', a'_{\alpha} \rangle && (6) \\ &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a'_{\alpha}, a_{\beta} \cdot b_i \rangle. \end{aligned}$$

This shows that two right A'' -module actions on $A^{(3)} = (A'')'$ and $A^{(3)} = (A')''$ are equal.

PROPOSITION 2.1 *Let A be a Banach algebra with following conditions*

- (i) A is Arens regular,
- (ii) the map $A \times A' \rightarrow A' ((a, a') \mapsto a \cdot a')$ is Arens regular.

Then two A'' -module actions on $A^{(3)} = (A')''$ and $A^{(3)} = (A'')'$ coincide.

Proof It is enough to prove that left module actions in (3) and (4) coincide. We begin with equation (3)

$$\begin{aligned}
 \langle a'' . a^{(3)}, b'' \rangle &= \lim_{\beta} \lim_{\alpha} \langle b'', a_{\beta} . a'_{\alpha} \rangle \\
 &= \lim_{\alpha} \lim_{\beta} \langle b'', a_{\beta} . a'_{\alpha} \rangle && \text{(by (ii))} \\
 &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a_{\beta} . a'_{\alpha}, b_i \rangle \\
 &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a'_{\alpha}, b_i . a_{\beta} \rangle && \text{(by formula (1))} \\
 &= \lim_{\alpha} \lim_i \lim_{\beta} \langle a'_{\alpha}, b_i . a_{\beta} \rangle && \text{(by (i))}
 \end{aligned}$$

this proves the equality of (3) and (4). ■

Now for $n = 2$ we consider two $A^{(4)}$ -module actions on $A^{(5)} = ((A')'')''$ and $A^{(5)} = ((A'')'')'$. Let $a^{(5)} = w^* - \lim_{\alpha} \widehat{a_{\alpha}^{(3)}} \in A^{(5)}$ and $a^{(4)} = w^* - \lim_{\beta} \widehat{a_{\beta}''}$, $b^{(4)} = w^* - \lim_i \widehat{b_i''}$ in $A^{(4)}$ where $(a_{\alpha}^{(3)})$ and $(a'_{\alpha}), (b_i'')$ are bounded nets in $A^{(3)}$ and A'' , respectively. For left $A^{(4)}$ -module action on $A^{(5)} = ((A')'')''$ we have

$$\begin{aligned}
 \langle a^{(4)} . a^{(5)}, b^{(4)} \rangle &= \lim_{\beta} \lim_{\alpha} \langle b^{(4)}, a_{\beta}'' . a_{\alpha}^{(3)} \rangle && \text{(by formula (2))} \\
 &= \lim_{\beta} \lim_{\alpha} \lim_i \langle a_{\beta}'' . a_{\alpha}^{(3)}, b_i'' \rangle && (7)
 \end{aligned}$$

and for left $A^{(4)}$ -module action on $A^{(5)} = ((A'')'')'$ we have

$$\begin{aligned}
 \langle a^{(4)} . a^{(5)}, b^{(4)} \rangle &= \langle a^{(5)}, b^{(4)} \square a^{(4)} \rangle && \text{(by formula (1))} \\
 &= \lim_{\alpha} \langle b^{(4)} \square a^{(4)}, a_{\alpha}^{(3)} \rangle && (8) \\
 &= \lim_{\alpha} \lim_i \lim_{\beta} \langle a_{\alpha}^{(3)}, b_i'' \square a_{\beta}'' \rangle.
 \end{aligned}$$

For right $A^{(4)}$ -module action on $A^{(5)} = ((A')'')''$ we have

$$\begin{aligned}
 \langle a^{(5)} . a^{(4)}, b^{(4)} \rangle &= \lim_{\alpha} \lim_{\beta} \langle b^{(4)}, a_{\alpha}^{(3)} . a_{\beta}'' \rangle && \text{(by formula (2))} \\
 &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a_{\alpha}^{(3)} . a_{\beta}'' , b_i'' \rangle, && (9)
 \end{aligned}$$

and for right $A^{(4)}$ -module action on $A^{(5)} = ((A'')'')'$

$$\begin{aligned}
 \langle a^{(5)} . a^{(4)}, b^{(4)} \rangle &= \langle a^{(5)}, a^{(4)} \square b^{(4)} \rangle && \text{(by formula (1))} \\
 &= \lim_{\alpha} \langle a^{(4)} \square b^{(4)}, a_{\alpha}^{(3)} \rangle && (10) \\
 &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a_{\alpha}^{(3)}, a_{\beta}'' \square b_i'' \rangle.
 \end{aligned}$$

We need the equality of two A'' -module actions on $A^{(3)} = (A'')'$ and $A^{(3)} = (A')''$ to prove the equality of above $A^{(4)}$ -module actions on $A^{(5)}$, so we need the following lemma whose proof is straightforward.

LEMMA 2.2 Let A be a Banach algebra with following conditions

- (i) A'' is Arens regular,
- (ii) the map $A'' \times A''' \rightarrow A'''$ $((a'', a^{(3)}) \mapsto a'' \cdot a^{(3)})$ is Arens regular.

Then the conditions of Proposition 2.1 hold.

PROPOSITION 2.3 Let A be a Banach algebra with conditions

- (i) A'' is Arens regular,
- (ii) the map $A'' \times A''' \rightarrow A'''$ $((a'', a^{(3)}) \mapsto a'' \cdot a^{(3)})$ is Arens regular.

Then two $A^{(4)}$ -module actions on $A^{(5)} = ((A')'')''$ and $A^{(5)} = ((A'')'')'$ coincide.

Proof By Lemma 2.2 the conditions of Proposition 2.1 hold, so two A'' -module actions on $A^{(3)} = (A')''$ and $A^{(3)} = (A'')'$ are equal. We begin with equality (7)

$$\begin{aligned} \langle a^{(4)} \cdot a^{(5)}, b^{(4)} \rangle &= \lim_{\beta} \lim_{\alpha} \langle b^{(4)}, a''_{\beta} \cdot a^{(3)}_{\alpha} \rangle \\ &= \lim_{\alpha} \lim_{\beta} \langle b^{(4)}, a''_{\beta} \cdot a^{(3)}_{\alpha} \rangle && \text{(by (ii) of Lemma 2.2)} \\ &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a''_{\beta} \cdot a^{(3)}_{\alpha}, b''_i \rangle \\ &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a^{(3)}_{\alpha}, b''_i \square a''_{\beta} \rangle && \text{(by Proposition 2.1)} \\ &= \lim_{\alpha} \lim_i \lim_{\beta} \langle a^{(3)}_{\alpha}, b''_i \square a''_{\beta} \rangle, && \text{(by (i) of Lemma 2.2)} \end{aligned}$$

this proves the equality of (7) and (8). For equality of right-module actions, we continue equality 9

$$\begin{aligned} \langle a^{(5)} \cdot a^{(4)}, b^{(4)} \rangle &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a^{(3)}_{\alpha} \cdot a''_{\beta}, b''_i \rangle \\ &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a^{(3)}_{\alpha}, a''_{\beta} \square b''_i \rangle && \text{(by Proposition 2.1)} \end{aligned}$$

and this proves the equality of (9) and (10). ■

Now suppose that $n = 3$, we consider two $A^{(6)}$ -module actions on $A^{(7)} = (((A')'')'')''$ and $A^{(7)} = (((A'')'')'')'$. Let $a^{(7)} = w^* - \widehat{\lim_{\alpha} a^{(5)}_{\alpha}} \in A^{(7)}$ and $a^{(6)} = w^* - \widehat{\lim_{\beta} a^{(4)}_{\beta}}$, $b^{(6)} = w^* - \widehat{\lim_i b^{(4)}_i} \in A^{(6)}$ where $(a^{(5)}_{\alpha})$ and $(a^{(4)}_{\beta}), (b^{(4)}_i)$ are bounded nets in $A^{(5)}$ and $A^{(4)}$, respectively. For left $A^{(6)}$ -module action on $A^{(7)} = (((A')'')'')''$ we can write

$$\langle a^{(6)} \cdot a^{(7)}, b^{(6)} \rangle = \lim_{\beta} \lim_{\alpha} \lim_i \langle a^{(4)}_{\beta} \cdot a^{(5)}_{\alpha}, b^{(4)}_i \rangle, \quad (11)$$

and for left $A^{(6)}$ -module action on $A^{(7)} = (((A'')''))'$ we can write

$$\langle a^{(6)}.a^{(7)}, b^{(6)} \rangle = \lim_{\alpha} \lim_i \lim_{\beta} \langle a_{\alpha}^{(5)}, b_i^{(4)} \square a_{\beta}^{(4)} \rangle. \tag{12}$$

For right $A^{(6)}$ -module action on $A^{(7)} = (((A')''))''$ we can write

$$\langle a^{(7)}.a^{(6)}, b^{(6)} \rangle = \lim_{\alpha} \lim_{\beta} \lim_i \langle a_{\alpha}^{(5)}.a_{\beta}^{(4)}, b_i^{(4)} \rangle, \tag{13}$$

and for right $A^{(6)}$ -module action on $A^{(7)} = (((A'')''))'$ we can write

$$\langle a^{(7)}.a^{(6)}, b^{(6)} \rangle = \lim_{\alpha} \lim_{\beta} \lim_i \langle a_{\alpha}^{(5)}, a_{\beta}^{(4)} \square b_i^{(4)} \rangle. \tag{14}$$

We need the equality of two A'' -module actions on $A^{(5)} = ((A'')'')$ and $A^{(5)} = ((A')'')$ to prove the equality of above $A^{(6)}$ -module actions on $A^{(7)}$, so we need the following Lemma that is similar to Lemma 2.2.

LEMMA 2.4 *Let A be a Banach algebra with following conditions*

- (i) $A^{(4)}$ is Arens regular,
- (ii) the map $A^{(4)} \times A^{(5)} \rightarrow A^{(5)} ((a^{(4)}, a^{(5)}) \mapsto a^{(4)}.a^{(5)})$ is Arens regular.

Then the conditions of Proposition 2.3 hold.

PROPOSITION 2.5 *Let A be a Banach algebra with conditions*

- (i) $A^{(4)}$ is Arens regular,
- (ii) the map $A^{(4)} \times A^{(5)} \rightarrow A^{(5)} ((a^{(4)}, a^{(5)}) \mapsto a^{(4)}.a^{(5)})$ is Arens regular.

Then two $A^{(6)}$ -module actions on $A^{(7)} = (((A'')''))'$ and $A^{(7)} = (((A')''))''$ coincide.

Proof By Lemma 2.4 the conditions of Proposition 2.3 hold, so two $A^{(4)}$ -module actions on $A^{(5)} = ((A')'')$ and $A^{(5)} = ((A'')'')$ are equal. We begin with equality (11)

$$\begin{aligned} \langle a^{(6)}.a^{(7)}, b^{(6)} \rangle &= \lim_{\beta} \lim_{\alpha} \langle b^{(6)}, a_{\beta}^{(4)}.a_{\alpha}^{(5)} \rangle \\ &= \lim_{\alpha} \lim_{\beta} \langle b^{(6)}, a_{\beta}^{(4)}.a_{\alpha}^{(5)} \rangle && \text{(by (ii) of Lemma 2.4)} \\ &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a_{\beta}^{(4)}.a_{\alpha}^{(5)}, b_i^{(4)} \rangle \\ &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a_{\alpha}^{(5)}, b_i^{(4)} \square a_{\beta}^{(4)} \rangle && \text{(by Proposition 2.3)} \\ &= \lim_{\alpha} \lim_i \lim_{\beta} \langle a_{\alpha}^{(5)}, b_i^{(4)} \square a_{\beta}^{(4)} \rangle, && \text{(by (i) of Lemma 2.4)} \end{aligned}$$

this prove the equality of (11) and (12). For equality of right-module actions, we

continue equality (13)

$$\begin{aligned} \langle a^{(7)}.a^{(6)}, b^{(6)} \rangle &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a_{\alpha}^{(5)}.a_{\beta}^{(4)}, b_i^{(4)} \rangle \\ &= \lim_{\alpha} \lim_{\beta} \lim_i \langle a_{\alpha}^{(5)}, a_{\beta}^{(4)} \square b_i^{(4)} \rangle \quad (\text{by Proposition 2.3}) \end{aligned}$$

and this proves the equality of (13) and (14). ■

Now by induction process we have the following extended result.

PROPOSITION 2.6 *Let A be a Banach algebra with following conditions for some $n \geq 1$*

- (i) A^{2n-2} is Arens regular,
- (ii) the map $A^{(2n-2)} \times A^{(2n-1)} \rightarrow A^{(2n-1)} ((a, f) \mapsto a.f)$ is Arens regular.

Then two $A^{(2n)}$ -module actions on $A^{(2n+1)} = (((A'')'' \cdots)'')$ and $A^{(2n+1)} = (((A')'' \cdots)''''$ coincide.

3. Main results

In this section we consider the transposes $D'', D^{(4)}, \dots, D^{(2n)}$ of a continuous derivation $D : A \rightarrow A'$. We know by Theorem 1.1 that the following maps will be continuous derivations

$$\begin{aligned} D'' &: A'' \longrightarrow A^{(3)} = (A')'' \\ D^{(4)} &: A^{(4)} = (A'')'' \longrightarrow A^{(5)} = ((A')'')'' \\ D^{(6)} &: A^{(6)} = ((A'')'')'' \longrightarrow A^{(7)} = (((A')'')'')'' \\ &\vdots \\ D^{(2n)} &: A^{(2n)} = (((A'')'') \cdots)'' \longrightarrow A^{(2n+1)} = (((A')'')'') \cdots)'' \end{aligned}$$

PROPOSITION 3.1 *Let A be a Banach algebra with hypothesis of Proposition 2.1. If the second transpose D'' of continuous derivation $D : A \rightarrow A'$ is inner, then D is inner.*

Proof Let $D : A \rightarrow A'$ be a derivation, then by Theorem 1.1 and Proposition 2.1, $D'' : A'' \rightarrow A^{(3)} = (A')'' = (A'')'$ is also a derivation. Since D'' is inner, there exists $a'' \in A''$ such that $D''(a'') = a'' \cdot a^{(3)} - a^{(3)} \cdot a''$, ($a^{(3)} \in A^{(3)}$). Let $a' = \iota^*(a^{(3)})$, where $\iota : A \rightarrow A''$ is the natural map. Then for each $a, b \in A$ we can write

$$\begin{aligned}
 \langle D(a), b \rangle &= \langle D''(\widehat{a}), \widehat{b} \rangle \\
 &= \langle \widehat{a}.a^{(3)} - a^{(3)}.\widehat{a}, \widehat{b} \rangle \\
 &= \langle a^{(3)}, \widehat{b}\square\widehat{a} - \widehat{a}\square\widehat{b} \rangle \quad (\text{by Proposition 2.1}) \\
 &= \langle a^{(3)}, \widehat{b.a - a.b} \rangle \\
 &= \langle a^{(3)}, \iota(b.a - a.b) \rangle \\
 &= \langle \iota^*(a^{(3)}), b.a - a.b \rangle \\
 &= \langle a', b.a - a.b \rangle \\
 &= \langle a.a' - a'.a, b \rangle,
 \end{aligned}$$

hence $D(a) = a.a' - a'.a$ and so D is inner. ■

PROPOSITION 3.2 *Let A be a Banach algebra with hypothesis of Proposition 2.3. If the fourth transpose $D^{(4)}$ of continuous derivation $D : A \rightarrow A'$ is inner, then D is inner.*

Proof Let $D : A \rightarrow A'$ be a derivation, then by Theorem 1.1 and Proposition 2.3, $D^{(4)} : ((A'')''') \rightarrow (((A')''')''')' = (((A'')''')''')'$ is also a derivation. Since $D^{(4)}$ is inner, there exists $a^{(4)} \in A^{(4)}$ such that $D^{(4)}(a^{(4)}) = a^{(4)}.a^{(5)} - a^{(5)}.a^{(4)}$, ($a^{(5)} \in A^{(5)}$). Let $a' = \iota^* \circ \iota^{***}(a^{(5)})$, where $\iota : A \rightarrow A''$ is the natural map. Then for each $a, b \in A$ we can write

$$\begin{aligned}
 \langle D(a), b \rangle &= \langle D''(\widehat{a}), \widehat{b} \rangle \\
 &= \langle \widehat{D''(\widehat{a})}, \widehat{\widehat{b}} \rangle \\
 &= \langle D^{(4)}(\widehat{\widehat{a}}), \widehat{\widehat{b}} \rangle \\
 &= \langle \widehat{\widehat{a}}.a^{(5)} - a^{(5)}.\widehat{\widehat{a}}, \widehat{\widehat{b}} \rangle \\
 &= \langle a^{(5)}, \widehat{\widehat{b}\square\widehat{a} - \widehat{\widehat{a}}\square\widehat{b}} \rangle \quad (\text{by Proposition 2.3}) \\
 &= \langle a^{(5)}, \widehat{b\square\widehat{a} - \widehat{a}\square\widehat{b}} \rangle \\
 &= \langle a^{(5)}, \iota^{**}(\widehat{b\square\widehat{a} - \widehat{a}\square\widehat{b}}) \rangle \\
 &= \langle \iota^{***}(a^{(5)}), b.a - a.b \rangle \\
 &= \langle \iota^{***}(a^{(5)}), \iota(b.a - a.b) \rangle \\
 &= \langle \iota^* \circ \iota^{***}(a^{(5)}), b.a - a.b \rangle \\
 &= \langle a', b.a - a.b \rangle \\
 &= \langle a.a' - a'.a, b \rangle,
 \end{aligned}$$

hence $D(a) = a.a' - a'.a$ and so D is inner. ■

PROPOSITION 3.3 *Let A be a Banach algebra with hypothesis of Proposition 2.5. If the sixth transpose $D^{(6)}$ of continuous derivation $D : A \rightarrow A'$ is inner, then D is inner.*

Proof Let $D : A \rightarrow A'$ be a derivation, then by Theorem 1.1 and Proposition 2.3, $D^{(6)} : A^{(6)} = ((A'')''')'' \rightarrow (((A')''')''')'' = (((A'')''')''')'' = A^{(7)}$ is also a derivation. Since $D^{(6)}$ is inner, there exists $a^{(6)} \in A^{(6)}$ such that $D^{(6)}(a^{(6)}) = a^{(6)}.a^{(7)} - a^{(7)}.a^{(6)}$, ($a^{(7)} \in A^{(7)}$). Let $a' = \iota^* \circ \iota^{***} \circ \iota^{*****}(a^{(7)})$, where $\iota : A \rightarrow A''$ is the

natural map. Then for each $a, b \in A$ we can write

$$\begin{aligned}
 \langle D(a), b \rangle &= \langle D''(\widehat{a}), \widehat{b} \rangle \\
 &= \langle \widehat{D''(\widehat{a})}, \widehat{\widehat{b}} \rangle \\
 &= \langle \widehat{D^{(4)}(\widehat{\widehat{a}})}, \widehat{\widehat{\widehat{b}}} \rangle \\
 &= \langle \widehat{D^{(6)}(\widehat{\widehat{\widehat{a}}})}, \widehat{\widehat{\widehat{\widehat{b}}}} \rangle \\
 &= \langle \widehat{\widehat{a}.a^{(7)}} - a^{(7)}. \widehat{\widehat{\widehat{a}}}, \widehat{\widehat{\widehat{b}}} \rangle \\
 &= \langle a^{(7)}, \widehat{\widehat{\widehat{b\Box a}}} - \widehat{\widehat{\widehat{a\Box b}}} \rangle && \text{(by Proposition 2.5)} \\
 &= \langle a^{(7)}, \widehat{\widehat{\widehat{b\Box a}}} - \widehat{\widehat{\widehat{a\Box b}}} \rangle \\
 &= \langle a^{(7)}, \iota^{****}(\widehat{\widehat{\widehat{b\Box a}}} - \widehat{\widehat{\widehat{a\Box b}}}) \rangle \\
 &= \langle \iota^{*****}(a^{(7)}), \iota^{**}(\widehat{\widehat{\widehat{b\Box a}}} - \widehat{\widehat{\widehat{a\Box b}}}) \rangle \\
 &= \langle \iota^{***} \circ \iota^{*****}(a^{(7)}), \widehat{\widehat{\widehat{b.a - a.b}}} \rangle \\
 &= \langle \iota^{***} \circ \iota^{*****}(a^{(7)}), \iota(b.a - a.b) \rangle \\
 &= \langle \iota^* \circ \iota^{***} \circ \iota^{*****}(a^{(7)}), b.a - a.b \rangle \\
 &= \langle a', b.a - a.b \rangle \\
 &= \langle a.a' - a'.a, b \rangle,
 \end{aligned}$$

hence $D(a) = a.a' - a'.a$ and so D is inner. \blacksquare

Using the similar reasoning as in the proof of previous lemmas we have the following proposition.

PROPOSITION 3.4 *Let A be a Banach algebra with hypothesis of Proposition 2.6. If the $(2n)$ -th transpose $D^{(2n)}$ of continuous derivation $D : A \rightarrow A'$ is inner, then D is inner.*

PROPOSITION 3.5 *Let A be a Banach algebra with hypothesis of Proposition 2.1. If A'' is weakly amenable, then A is weakly amenable.*

Proof Suppose that $D : A \rightarrow A'$ be a continuous derivation. Then $D'' : A'' \rightarrow A^{(3)} = (A')''$ is a continuous derivation by Theorem 1.1. But two A'' -module actions on $A^{(3)} = (A')''$ and $A^{(3)} = (A'')'$ are equal by Proposition 2.1, hence $D'' : A'' \rightarrow A^{(3)} = (A'')'$ is also a continuous derivation in which $A^{(3)} = (A'')'$ is considered as dual of A'' . Since A'' is weakly amenable, then D'' is inner. Therefore D is inner by Proposition 3.1. This completes the proof. \blacksquare

Using the same reasoning as in the proofs of previous propositions we have next results, so we omit the details in proofs.

PROPOSITION 3.6 *Let A be a Banach algebra with hypothesis of Proposition 2.3. If $A^{(4)}$ is weakly amenable, then A is weakly amenable.*

Proof This is a consequence of Proposition 3.2. \blacksquare

PROPOSITION 3.7 *Let A be a Banach algebra with hypothesis of Proposition 2.5. If $A^{(6)}$ is weakly amenable, then A is weakly amenable.*

Proof This is a consequence of Proposition 3.3. ■

Finally by Propositions 2.6 and 3.4 we have the following extended result.

PROPOSITION 3.8 *Let A be a Banach algebra with hypothesis of Proposition 2.6. If $A^{(2n)}$ is weakly amenable, then A is weakly amenable.*

Acknowledgements: The authors would like to thank the referee for carefully reading and giving some fruitful suggestions.

References

- [1] R. ARENS, The adjoint of a bilinear operation, Proc. Amer. Math. Soc. 2 (1951), 839-848.
- [2] W.G. BADE, P.C. CURTIS AND H.G. DALES,, Amenability and weak amenability for Bearling and Lipschitz algebra, Proc. London Math. Soc. , 55 (1987), no. 3, 359-377.
- [3] A. BODAGHI, M.ETTEFAGH, M.E. GORDJI AND A. MEDGHALCHI, Module structures on iterated duals of Banach algebras, An.st.Univ.Ovidius Constanta, 18(1) (2010) 63-80.
- [4] H.G. DALES, F. GHARAMANI, AND N. GRONBAEK, Derivations into iterated duals of Banach algebras, Studia Math, 128 (1998), no.1, 19-54.
- [5] H.G. DALES, Banach algebra and Automatic continuity, Oxford university Press, (2000).
- [6] M. ETTEFAGH, The third dual of a Banach algebra, Studia. Sci. Math. Hung, 45(1) (2008) 1-11.
- [7] B.E. JOHNSON, Cohomology in Banach algebras, Mem. Amer. Math. Soc, 127 (1972).
- [8] A. Medghalchi and T.Yazdanpanah, Problems concerning n-weak amenability of a Banach algebra, Czechoslovak Math. J, 55(130) (2005) 863-876.