

## On weakly $eR$ -open functions

M. Özkoç<sup>a\*</sup>, B. S. Ayhan<sup>a</sup>

<sup>a</sup>Department of Mathematics, Faculty of Science Muğla Sıtkı Koçman University,  
Menteşe-Muğla 48000 Turkey.

Received 28 March 2016; Revised 24 June 2016; Accepted 30 June 2016.

---

**Abstract.** The main goal of this paper is to introduce and study a new class of function via the notions of  $e$ - $\theta$ -open sets and  $e$ - $\theta$ -closure operator which are defined by Özkoç and Aslım [10] called weakly  $eR$ -open functions and  $e$ - $\theta$ -open functions. Moreover, we investigate not only some of their basic properties but also their relationships with other types of already existing topological functions.

© 2016 IAUCTB. All rights reserved.

---

**Keywords:**  $e$ -Closed set,  $e$ - $\theta$ -open set, weakly  $eR$ -open function,  $e$ - $\theta$ -open function.

**2010 AMS Subject Classification:** Primary 54C08; Secondary 54D05.

### 1. Introduction and Preliminaries

Throughout the present paper,  $X$  and  $Y$  always mean topological spaces on which no separation axioms are assumed unless explicitly stated. Let  $X$  be a topological space and  $A$  a subset of  $X$ . The closure and the interior of  $A$  are denoted by  $cl(A)$  and  $int(A)$ , respectively. The family of all closed sets of  $X$  is denoted  $C(X)$ . A subset  $A$  is said to be regular open [12] (resp. regular closed [12]) if  $A = int(cl(A))$  (resp.  $A = cl(int(A))$ ). A point  $x \in X$  is said to be  $\delta$ -cluster point [13] of  $A$  if  $int(cl(U)) \cap A \neq \emptyset$  for each open neighbourhood  $U$  of  $x$ . The set of all  $\delta$ -cluster points of  $A$  is called the  $\delta$ -closure [13] of  $A$  and is denoted by  $cl_\delta(A)$ . If  $A = cl_\delta(A)$ , then  $A$  is called  $\delta$ -closed [13], and the complement of a  $\delta$ -closed set is called  $\delta$ -open [13]. A subset  $A$  is called semiopen [5] (resp.  $b$ -open [1],  $e$ -open [4], preopen [7],  $\alpha$ -open [8]) if  $A \subset cl(int(A))$  (resp.  $A \subset cl(int(A)) \cup int(cl(A))$ ,  $A \subset cl(int_\delta(A)) \cup int(cl_\delta(A))$ ,  $A \subset int(cl(A))$ ,

---

\*Corresponding author.

E-mail address: murad.ozkoc@mu.edu.tr (M. Özkoç).

$A \subset \text{int}(cl(\text{int}(A)))$ ). The complement of a semiopen (resp.  $b$ -open,  $e$ -open, preopen,  $\alpha$ -open) set is called semiclosed [5](resp.  $b$ -closed [1],  $e$ -closed [4], preclosed [7],  $\alpha$ -closed [8]). The intersection of all  $e$ -closed sets of  $X$  containing  $A$  is called the  $e$ -closure [4] of  $A$  and is denoted by  $e-cl(A)$ . The union of all  $e$ -open sets of  $X$  contained in  $A$  is called the  $e$ -interior [4] of  $A$  and is denoted by  $e-int(A)$ . A subset  $A$  is said to be  $e$ -regular [10] if it is  $e$ -open and  $e$ -closed.

A point  $x$  of  $X$  is called a  $b$ - $\theta$ -cluster [11] ( $e$ - $\theta$ -cluster [10],  $\theta$ -cluster [13]) point of  $A$  if  $bcl(U) \cap A \neq \emptyset$  ( $e-cl(U) \cap A \neq \emptyset$ ,  $cl(U) \cap A \neq \emptyset$ ) for every  $b$ -open ( $e$ -open, open) set  $U$  of  $X$  containing  $x$ , respectively. The set of all  $b$ - $\theta$ -cluster ( $e$ - $\theta$ -cluster,  $\theta$ -cluster) points of  $A$  is called the  $b$ - $\theta$ -closure [11] ( $e$ - $\theta$ -closure [10],  $\theta$ -closure [13]) of  $A$  and is denoted by  $bcl_\theta(A)$  ( $e-cl_\theta(A)$ ,  $cl_\theta(A)$ ), respectively. A subset  $A$  is said to be  $b$ - $\theta$ -closed [11] ( $e$ - $\theta$ -closed [10],  $\theta$ -closed [13]) if  $A = bcl_\theta(A)$  ( $A = e-cl_\theta(A)$ ,  $A = cl_\theta(A)$ ), respectively. The complement of a  $b$ - $\theta$ -closed ( $e$ - $\theta$ -closed,  $\theta$ -closed) set is called a  $b$ - $\theta$ -open [11] ( $e$ - $\theta$ -open [10],  $\theta$ -open [13]) set. A point  $x$  of  $X$  said to be a  $b$ - $\theta$ -interior [11] ( $e$ - $\theta$ -interior [10],  $\theta$ -interior [13]) point of a subset  $A$ , denoted by  $bint_\theta(A)$  ( $e-int_\theta(A)$ ,  $int_\theta(A)$ ), if there exists a  $b$ -regular ( $e$ -regular, regular) set  $U$  of  $X$  containing  $x$  such that  $U \subset A$ , respectively. The family of all  $e$ -open (resp.  $e$ -closed,  $e$ -regular,  $b$ - $\theta$ -open,  $e$ - $\theta$ -open,  $b$ - $\theta$ -closed,  $e$ - $\theta$ -closed) subsets of  $X$  is denoted by  $eO(X)$  (resp.  $eC(X)$ ,  $eR(X)$ ,  $B\theta O(X)$ ,  $e\theta O(X)$ ,  $B\theta C(X)$ ,  $e\theta C(X)$ ). The family of all  $e$ -open ( $e$ -closed,  $e$ -regular,  $b$ - $\theta$ -open,  $e$ - $\theta$ -open,  $b$ - $\theta$ -closed,  $e$ - $\theta$ -closed) sets of  $X$  containing a point  $x$  of  $X$  is denoted by  $eO(X, x)$  (resp.  $eC(X, x)$ ,  $eR(X, x)$ ,  $B\theta O(X, x)$ ,  $e\theta O(X, x)$ ,  $B\theta C(X, x)$ ,  $e\theta C(X, x)$ ). Also it is noted in [10] that

$$e\text{-regular} \Rightarrow e\text{-}\theta\text{-open} \Rightarrow e\text{-open}.$$

We shall use the well-known accepted language almost in the whole of the article.

**Definition 1.1** A function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is called:

- (a) contra  $e$ - $\theta$ -open if  $f(U)$  is  $e$ - $\theta$ -closed in  $Y$  for each open set  $U$  of  $X$ .
- (b) contra  $e$ - $\theta$ -closed if  $f(U)$  is  $e$ - $\theta$ -open in  $Y$  for each closed set  $U$  of  $X$ .
- (c) strongly continuous [6] if for every subset  $A$  of  $X$ ,  $f(cl(A)) \subset f(A)$ .
- (d) weakly  $BR$ -open [2] if  $f(U) \subset bint_\theta(f(cl(U)))$  for each open set  $U$  of  $X$ .

## 2. Weakly $eR$ -open Functions

In this section, we define the concept of weakly  $eR$ -open and investigate some basic properties of them.

**Definition 2.1** A function  $f : X \rightarrow Y$  is said to be weakly  $eR$ -open if  $f(U) \subset e-int_\theta(f(cl(U)))$  for each open set  $U$  of  $X$ .

**Definition 2.2** A function  $f : X \rightarrow Y$  is said to be  $e$ - $\theta$ -open if  $f(U)$  is  $e$ - $\theta$ -open in  $Y$  for each open set  $U$  of  $X$ .

It is clear to see that every  $e$ - $\theta$ -open function is a weakly  $eR$ -open. However, a weakly  $eR$ -open function need not be  $e$ - $\theta$ -open as shown by the following example.

**Example 2.3** Let  $X = \{a, b, c, d\}$  and

$$\tau = \{\emptyset, X, \{a, d\}\} \quad \text{and} \quad \sigma = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, c\}, \{a, b, c\}, \{a, b, d\}\}.$$

The identity function  $f : (X, \tau) \rightarrow (X, \sigma)$  is weakly  $eR$ -open, but it is not  $e$ - $\theta$ -open.

The notions of weakly  $eR$ -open function and weakly  $BR$ -open function are independent as shown by the following examples.

**Example 2.4** Let  $X = \{a, b, c, d, e\}$  and  $\tau = \{\emptyset, X, \{a\}, \{c\}, \{a, c\}, \{c, d\}, \{a, c, d\}\}$ . The identity function  $f : (X, \tau) \rightarrow (X, \tau)$  is weakly  $eR$ -open, but it is not weakly  $BR$ -open.

**Example 2.5** Let  $X = \{a, b, c, d\}$  and  $\tau = \{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{a, c\}, \{a, b, c\}, \{a, b, d\}\}$ .  $f = \{(a, d), (b, c), (c, b), (d, d)\}$  is weakly  $BR$ -open, but it is not weakly  $eR$ -open.

**Lemma 2.6** [10] Let  $A$  be a subset of a space  $X$ . Then:

- (1)  $e-cl_\theta(A) = \cap\{V | (A \subset V)(V \in eR(X))\}$ .
- (2)  $x \in e-cl_\theta(A)$  iff  $A \cap U \neq \emptyset$  for each  $e$ -regular set  $U$  of  $X$  containing  $x$ .
- (3)  $e-cl_\theta(A)$  is  $e$ - $\theta$ -closed.
- (4) Any intersections of  $e$ - $\theta$ -closed sets is  $e$ - $\theta$ -closed and any union of  $e$ - $\theta$ -open sets is  $e$ - $\theta$ -open.
- (5)  $A$  is  $e$ - $\theta$ -open in  $X$  if and only if for each  $x \in A$  there exists an  $e$ -regular set  $U$  containing  $x$  such that  $x \in U \subset A$ .

**Theorem 2.7** Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a function. Then the following statements are equivalent:

- (a)  $f$  is weakly  $eR$ -open,
- (b)  $f(int_\theta(A)) \subset e-int_\theta(f(A))$  for every subset  $A$  of  $X$ ,
- (c)  $int_\theta(f^{-1}(B)) \subset f^{-1}(e-int_\theta(B))$  for every subset  $B$  of  $Y$ ,
- (d)  $f^{-1}(e-cl_\theta(B)) \subset cl_\theta(f^{-1}(B))$  for every subset  $B$  of  $Y$ ,
- (e)  $f(int(F)) \subset e-int_\theta(f(F))$  for each closed subset  $F$  of  $X$ ,
- (f)  $f(int(cl(U))) \subset e-int_\theta(f(cl(U)))$  for each open subset  $U$  of  $X$ ,
- (g)  $f(U) \subset e-int_\theta(f(cl(U)))$  for every regular open subset  $U$  of  $X$ ,
- (h)  $f(U) \subset e-int_\theta(f(cl(U)))$  for every  $\alpha$ -open subset  $U$  of  $X$ ,
- (i) For each  $x \in X$  and each open set  $U$  of  $X$  containing  $x$ , there exists an  $e$ - $\theta$ -open set  $V$  of  $Y$  containing  $f(x)$  such that  $V \subset f(cl(U))$ .

**Proof.** (a)  $\Rightarrow$  (b): Let  $A$  be any subset of  $X$  and  $x \in int_\theta(A)$ .

$$\begin{aligned} x \in int_\theta(A) &\Rightarrow (\exists U \in \mathcal{U}(x))(x \in U \subset cl(U) \subset A) \\ &\Rightarrow (\exists U \in \mathcal{U}(x))(f(x) \in f(U) \subset f(cl(U)) \subset f(A)) \Big\} \Rightarrow \\ &\hspace{15em} f \text{ is weakly } eR\text{-open} \\ &\Rightarrow f(U) \subset e-int_\theta(f(cl(U))) \subset e-int_\theta(f(A)) \\ &\Rightarrow f(x) \in e-int_\theta(f(A)) \\ &\Rightarrow x \in f^{-1}(e-int_\theta(f(A))). \end{aligned}$$

(b)  $\Rightarrow$  (c): Let  $B$  be any subset of  $Y$ .

$$\begin{aligned} B \subset Y \Rightarrow f^{-1}(B) \subset X \Big\} &\Rightarrow f(int_\theta(f^{-1}(B))) \subset e-int_\theta(f(f^{-1}(B))) \subset e-int_\theta(B) \\ (b) &\Rightarrow int_\theta(f^{-1}(B)) \subset f^{-1}(e-int_\theta(B)). \end{aligned}$$

(c)  $\Rightarrow$  (d): Let  $B$  be any subset of  $Y$ .

$$\left. \begin{aligned} B \subset Y \Rightarrow Y \setminus B \subset Y \\ (c) \end{aligned} \right\} \Rightarrow \begin{aligned} & \text{int}_\theta(f^{-1}(Y \setminus B)) \subset f^{-1}(e\text{-int}_\theta(Y \setminus B)) \\ & \Rightarrow \text{int}_\theta(X \setminus f^{-1}(B)) \subset f^{-1}(Y \setminus e\text{-cl}_\theta(B)) \\ & \Rightarrow X \setminus \text{cl}_\theta(f^{-1}(B)) \subset X \setminus f^{-1}(e\text{-cl}_\theta(B)) \\ & \Rightarrow f^{-1}(e\text{-cl}_\theta(B)) \subset \text{cl}_\theta(f^{-1}(B)). \end{aligned}$$

(d)  $\Rightarrow$  (e): Let  $F$  be any closed set of  $X$ .

$$\left. \begin{aligned} F \in C(X) \Rightarrow Y \setminus f(F) \subset Y \\ (d) \end{aligned} \right\} \Rightarrow \begin{aligned} & \Rightarrow f^{-1}(e\text{-cl}_\theta(Y \setminus f(F))) \subset \text{cl}_\theta(f^{-1}(Y \setminus f(F))) = \text{cl}_\theta(X \setminus f^{-1}(f(F))) \subset \text{cl}_\theta(X \setminus F) \\ & \Rightarrow f^{-1}(Y \setminus e\text{-int}_\theta(f(F))) \subset \text{cl}_\theta(X \setminus F) = X \setminus \text{int}_\theta(F) \\ & \Rightarrow X \setminus f^{-1}(e\text{-int}_\theta(f(F))) \subset X \setminus \text{int}_\theta(F) \end{aligned} \right\} \Rightarrow \begin{aligned} & f(\text{int}(F)) \subset e\text{-int}_\theta(f(F)). \\ & F \in C(X) \Rightarrow \text{int}_\theta(F) = \text{int}(F) \end{aligned}$$

(e)  $\Rightarrow$  (f), (f)  $\Rightarrow$  (g): Obvious.

(g)  $\Rightarrow$  (h): Let  $U$  be any  $\alpha$ -open set of  $X$ .

$$\left. \begin{aligned} U \in \alpha O(X) \Rightarrow (U \subset \text{int}(\text{cl}(\text{int}(U))))(\text{int}(\text{cl}(\text{int}(U))) \in RO(X)) \\ (g) \end{aligned} \right\} \Rightarrow \begin{aligned} & \Rightarrow f(U) \subset f(\text{int}(\text{cl}(\text{int}(U)))) \subset e\text{-int}_\theta(f(\text{cl}(\text{int}(\text{cl}(\text{int}(U)))))) \\ & = e\text{-int}_\theta(f(\text{cl}(\text{int}(U)))) \subset e\text{-int}_\theta(f(\text{cl}(U))). \end{aligned}$$

(h)  $\Rightarrow$  (i): Straightforward.

(i)  $\Rightarrow$  (a): Let  $U$  be an open set in  $X$  and  $y \in f(U)$ .

$$\left. \begin{aligned} (U \in \tau)(y \in f(U)) \\ (i) \end{aligned} \right\} \Rightarrow \left. \begin{aligned} (\exists V \in e\theta O(Y, y))(V \subset f(\text{cl}(U))) \\ y \in V \subset e\text{-int}_\theta(f(\text{cl}(U))) \end{aligned} \right\} \Rightarrow \begin{aligned} & \Rightarrow f(U) \subset e\text{-int}_\theta(f(\text{cl}(U))). \end{aligned}$$

■

**Theorem 2.8** Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a bijective function. Then the following statements are equivalent:

- (a)  $f$  is weakly  $eR$ -open,
- (b) For each  $x \in X$  and each open set  $U$  of  $X$  containing  $x$ , there exists an  $e$ -regular set  $V$  containing  $f(x)$  such that  $V \subset f(\text{cl}(U))$ ,
- (c)  $e\text{-cl}_\theta(f(\text{int}(\text{cl}(U)))) \subset f(\text{cl}(U))$  for each subset  $U$  of  $X$ ,
- (d)  $e\text{-cl}_\theta(f(\text{int}(F))) \subset f(F)$  for each regular closed subset  $F$  of  $X$ ,
- (e)  $e\text{-cl}_\theta(f(U)) \subset f(\text{cl}(U))$  for each open subset  $U$  of  $X$ ,
- (f)  $e\text{-cl}_\theta(f(U)) \subset f(\text{cl}(U))$  for each preopen subset  $U$  of  $X$ ,
- (g)  $f(U) \subset e\text{-int}_\theta(f(\text{cl}(U)))$  for each preopen subset  $U$  of  $X$ ,
- (h)  $f^{-1}(e\text{-cl}_\theta(B)) \subset \text{cl}_\theta(f^{-1}(B))$  for each subset  $B$  of  $Y$ ,
- (i)  $e\text{-cl}_\theta(f(U)) \subset f(\text{cl}_\theta(U))$  for each subset  $U$  of  $X$ ,
- (j)  $e\text{-cl}_\theta(f(\text{int}(\text{cl}_\theta(U)))) \subset f(\text{cl}_\theta(U))$  for each subset  $U$  of  $X$ .

**Proof.** (a)  $\Rightarrow$  (b): Let  $x \in X$  and  $U$  be any open subset of  $X$  containing  $x$ .

$$\left. \begin{aligned} x \in U \in \tau \\ (a) \end{aligned} \right\} \Rightarrow \left. \begin{aligned} f(x) \in f(U) \subset e\text{-int}_\theta(f(\text{cl}(U))) \in e\theta O(Y, f(x)) \\ \text{Lemma 2.6(5)} \end{aligned} \right\} \Rightarrow \begin{aligned} & \Rightarrow (\exists V \in eR(Y, f(x)))(V \subset e\text{-int}_\theta(f(\text{cl}(U))) \subset f(\text{cl}(U))). \end{aligned}$$

(b)  $\Rightarrow$  (c): Let  $x \in X$  and  $U \subset X$ .

$$\begin{aligned} f(x) \in Y \setminus f(\text{cl}(U)) = f(X \setminus \text{cl}(U)) &\Rightarrow x \in X \setminus \text{cl}(U) \\ &\Rightarrow (\exists G \in \mathcal{U}(x))(G \cap U = \emptyset) \\ &\Rightarrow (\exists G \in \mathcal{U}(x))(\text{cl}(G) \cap \text{int}(\text{cl}(U)) = \emptyset) \} \Rightarrow \\ &\hspace{15em} (b) \\ &\Rightarrow (\exists V \in eR(Y, f(x)))(V \subset f(\text{cl}(G))) \\ &\Rightarrow (\exists V \in eR(Y, f(x)))(V \cap f(\text{int}(\text{cl}(U))) = \emptyset) \\ &\Rightarrow f(x) \notin e\text{-cl}_\theta(f(\text{int}(\text{cl}(U)))) \\ &\Rightarrow f(x) \in X \setminus e\text{-cl}_\theta(f(\text{int}(\text{cl}(U)))). \end{aligned}$$

(c)  $\Rightarrow$  (d): Let  $F$  be any regular closed set of  $X$ .

$$\begin{aligned} F \in RC(X) \Rightarrow e\text{-cl}_\theta(f(\text{int}(F))) = e\text{-cl}_\theta(f(\text{int}(\text{cl}(\text{int}(F)))))) \} \Rightarrow \\ \hspace{15em} (c) \\ \Rightarrow e\text{-cl}_\theta(f(\text{int}(F))) \subset f(\text{cl}(\text{int}(F))) = f(F). \end{aligned}$$

(d)  $\Rightarrow$  (e): Let  $U$  be any open subset of  $X$ .

$$(U \in \tau)(\text{cl}(U) \in RC(X)) \} \Rightarrow e\text{-cl}_\theta(f(U)) \subset e\text{-cl}_\theta(f(\text{int}(\text{cl}(U)))) \subset f(\text{cl}(U)).$$

(d)

(e)  $\Rightarrow$  (f): Let  $U$  be any preopen subset of  $X$ .

$$\begin{aligned} U \in PO(X) \Rightarrow (U \subset \text{int}(\text{cl}(U)))(\text{int}(\text{cl}(U)) \in \tau) \} \Rightarrow \\ \hspace{15em} (e) \\ \Rightarrow e\text{-cl}_\theta(f(U)) \subset e\text{-cl}_\theta(f(\text{int}(\text{cl}(U)))) \subset f(\text{cl}(\text{int}(\text{cl}(U)))) \subset f(\text{cl}(U)). \end{aligned}$$

(f)  $\Rightarrow$  (g): Let  $U$  be any preopen subset of  $X$ .

$$\begin{aligned} U \in PO(X) \Rightarrow X \setminus \text{cl}(U) \in \tau \} \Rightarrow e\text{-cl}_\theta(f(X \setminus \text{cl}(U))) \subset f(\text{cl}(X \setminus \text{cl}(U))) \\ \hspace{15em} (f) \\ \Rightarrow e\text{-cl}_\theta(Y \setminus f(\text{cl}(U))) \subset f(X \setminus \text{int}(\text{cl}(U))) = Y \setminus f(\text{int}(\text{cl}(U))) \\ \Rightarrow Y \setminus e\text{-int}_\theta(f(\text{cl}(U))) \subset Y \setminus f(\text{int}(\text{cl}(U))) \\ \Rightarrow f(U) \subset f(\text{int}(\text{cl}(U))) \subset e\text{-int}_\theta(f(\text{cl}(U))). \end{aligned}$$

(g)  $\Rightarrow$  (h): Straightforward.

(h)  $\Rightarrow$  (i): Let  $U \subset X$ .

$$\begin{aligned} U \subset X \Rightarrow f(U) \subset Y \} \Rightarrow f^{-1}(e\text{-cl}_\theta(f(U))) \subset \text{cl}_\theta(f^{-1}(f(U))) = \text{cl}_\theta(U) \\ \hspace{15em} (h) \\ \Rightarrow e\text{-cl}_\theta(f(U)) \subset f(\text{cl}_\theta(U)). \end{aligned}$$

(i)  $\Rightarrow$  (j): Let  $U \subset X$ .

$$\begin{aligned} U \subset X \Rightarrow \text{cl}_\theta(U) \in C(X) \Rightarrow \text{int}(\text{cl}_\theta(U)) \subset X \} \Rightarrow \\ \hspace{15em} (i) \\ \Rightarrow e\text{-cl}_\theta(f(\text{int}(\text{cl}_\theta(U)))) \subset f(\text{cl}_\theta(\text{int}(\text{cl}_\theta(U)))) = f(\text{cl}(\text{int}(\text{cl}_\theta(U)))) \subset f(\text{cl}_\theta(U)). \end{aligned}$$

(j)  $\Rightarrow$  (a): Straightforward. ■

**Theorem 2.9** If  $X$  is a regular space and  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a bijective function, then the following statements are equivalent:

- (a)  $f$  is weakly  $eR$ -open.
- (b) For each  $\theta$ -open set  $A$  in  $X$ ,  $f(A)$  is  $e$ - $\theta$ -open in  $Y$ .

(c) For any set  $B$  of  $Y$  and any  $\theta$ -closed set  $A$  in  $X$  containing  $f^{-1}(B)$ , there exists an  $e$ - $\theta$ -closed set  $F$  in  $Y$  containing  $B$  such that  $f^{-1}(F) \subset A$ .

**Proof.** (a)  $\Rightarrow$  (b): Let  $A$  be a  $\theta$ -open set in  $X$ .

$$\begin{aligned} A \in \theta O(X) \Rightarrow Y \setminus f(A) \subset Y \} &\Rightarrow f^{-1}(e-cl_{\theta}(Y \setminus f(A))) \subset cl_{\theta}(f^{-1}(Y \setminus f(A))) \\ (a)(\text{Theorem 2.7}(d)) \} &\Rightarrow X \setminus f^{-1}(e-int_{\theta}(f(A))) \subset cl_{\theta}(X \setminus A) = X \setminus A \\ &\Rightarrow A \subset f^{-1}(e-int_{\theta}(f(A))) \\ &\Rightarrow f(A) \subset e-int_{\theta}(f(A)). \end{aligned}$$

(b)  $\Rightarrow$  (c): Let  $B$  be any set in  $Y$  and  $A$  be a  $\theta$ -closed set in  $X$  such that  $f^{-1}(B) \subset A$ .

$$\begin{aligned} (B \subset Y)(A \in \theta C(X))(f^{-1}(B) \subset A) &\Rightarrow (X \setminus A \in \theta O(X))(B \subset Y \setminus f(X \setminus A)) \} \Rightarrow \\ (b) \} &\Rightarrow (f(X \setminus A) \in e\theta O(Y))(B \subset Y \setminus f(X \setminus A)) \} \\ F := Y \setminus f(X \setminus A) \} &\Rightarrow \\ \Rightarrow (F \in e\theta C(X))(B \subset Y)(f^{-1}(F) = f^{-1}(Y \setminus f(X \setminus A))) &= f^{-1}(f(A)) = A. \end{aligned}$$

(c)  $\Rightarrow$  (a) : Let  $B$  be any set in  $Y$ .

$$\begin{aligned} (B \subset Y)(f^{-1}(B) \subset cl_{\theta}(f^{-1}(B))) \} &\Rightarrow \\ X \text{ is regular} \Rightarrow cl_{\theta}(f^{-1}(B)) \in \theta C(X) \} &\Rightarrow \\ (c) \} &\Rightarrow (\exists F \in e\theta C(Y))(B \subset F)(f^{-1}(F) \subset cl_{\theta}(f^{-1}(B))) \\ \Rightarrow (\exists F \in e\theta C(Y))(B \subset F)(f^{-1}(e-cl_{\theta}(B)) \subset f^{-1}(F) \subset cl_{\theta}(f^{-1}(B))). & \end{aligned}$$

Then from Theorem 2.8(h)  $f$  is weakly  $eR$ -open. ■

**Theorem 2.10** If  $X$  is a regular space and  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a bijective function, then the following statements are equivalent:

- (a)  $f$  is weakly  $eR$ -open.
- (b)  $f$  is  $e$ - $\theta$ -open.
- (c) For each  $x \in X$  and each open set  $U$  of  $X$  containing  $x$ , there exists an  $e$ -open set  $V$  of  $Y$  containing  $f(x)$  such that  $e-cl(V) \subset f(U)$ .

**Proof.** (a)  $\Rightarrow$  (b): Let  $W$  be a nonempty open subset of  $X$ .

$$\begin{aligned} x \in W \in \tau \} &\Rightarrow (\exists U_x \in \mathcal{U}(x))(cl(U_x) \subset W) \\ X \text{ is regular} \} &\Rightarrow W = \cup\{U_x|x \in W\} = \cup\{cl(U_x)|x \in W\} \\ \Rightarrow f(W) = \cup\{f(U_x)|x \in W\} \} &\Rightarrow f(W) = \cup\{f(U_x)|x \in W\} \\ f \text{ is weakly } eR\text{-open} \} &\Rightarrow \cup\{e-int_{\theta}(f(cl(U_x)))|x \in W\} \\ &\subset e-int_{\theta}(\cup\{f(cl(U_x))|x \in W\}) \} \Rightarrow \\ & f \text{ is bijective} \} \\ \Rightarrow f(W) \subset e-int_{\theta}(f(\cup\{cl(U_x)|x \in W\})) = e-int_{\theta}(f(W)) \} &\Rightarrow \\ e-int_{\theta}(f(W)) \subset f(W) \} &\Rightarrow \\ \Rightarrow e-int_{\theta}(f(W)) = f(W) & \\ \Rightarrow f(W) \in e\theta O(Y). & \end{aligned}$$

(b)  $\Rightarrow$  (c) and (c)  $\Rightarrow$  (a): Straightforward. ■

**Theorem 2.11** If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is weakly  $eR$ -open and strongly continuous, then

$f$  is  $e$ - $\theta$ -open.

**Proof.** Let  $U$  be any open subset of  $X$ .

$$\left. \begin{array}{l} U \in \tau \\ f \text{ is weakly } eR\text{-open} \end{array} \right\} \Rightarrow \left. \begin{array}{l} f(U) \subset e\text{-int}_\theta(f(\text{cl}(U))) \\ f \text{ is strongly continuous} \end{array} \right\} \Rightarrow \\ \Rightarrow f(U) \subset e\text{-int}_\theta(f(\text{cl}(U))) \subset e\text{-int}_\theta(f(U)). \quad \blacksquare$$

The following example shows that strong continuity is not decomposition of  $e$ - $\theta$ -openness. Namely, an  $e$ - $\theta$ -open function need not be strongly continuous.

**Example 2.12** Let  $X = \{a, b\}$  and  $\tau$  be the indiscrete topology for  $X$ . Then the identity function  $f : (X, \tau) \rightarrow (X, \tau)$  is an  $e$ - $\theta$ -open function but it is not strongly continuous.

**Theorem 2.13** If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is contra  $e$ - $\theta$ -closed, then  $f$  is a weakly  $eR$ -open function.

**Proof.** Let  $U$  be any open subset of  $X$ .

$$\left. \begin{array}{l} U \in \tau \Rightarrow \text{cl}(U) \in C(X) \\ f \text{ is contra } e\text{-}\theta\text{-closed} \end{array} \right\} \Rightarrow f(\text{cl}(U)) \in e\theta O(Y) \\ \Rightarrow f(U) \subset f(\text{cl}(U)) = e\text{-int}_\theta(f(\text{cl}(U))). \quad \blacksquare$$

**Theorem 2.14** If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is bijective contra  $e$ - $\theta$ -open, then  $f$  is a weakly  $eR$ -open function.

**Proof.** Let  $U$  be any open subset of  $X$ .

$$\left. \begin{array}{l} U \in \tau \\ f \text{ is contra } e\text{-}\theta\text{-open} \end{array} \right\} \Rightarrow f(U) \in e\theta C(Y) \Rightarrow e\text{-cl}_\theta(f(U)) = f(U) \subset f(\text{cl}(U)).$$

Then from Theorem 2.8(e)  $f$  is weakly  $eR$ -open. ■

**Theorem 2.15** Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a bijective function. If  $f(\text{cl}_\theta(U))$  is  $e$ - $\theta$ -closed in  $Y$  for every subset  $U$  of  $X$ , then  $f$  is weakly  $eR$ -open.

**Proof.** Let  $U$  be a subset of  $X$ .

$$(U \subset X)(f(\text{cl}_\theta(U)) \in e\theta C(Y)) \Rightarrow e\text{-cl}_\theta(f(U)) \subset e\text{-cl}_\theta(f(\text{cl}_\theta(U))) = f(\text{cl}_\theta(U)).$$

Then from Theorem 2.8(i)  $f$  is weakly  $eR$ -open. ■

**Definition 2.16** A function  $f : X \rightarrow Y$  is called complementary weakly  $eR$ -open (briefly c.w. $eR$ -o) if for each open set  $U$  of  $X$ ,  $f(\text{Fr}(U))$  is  $e$ - $\theta$ -closed in  $Y$ , where  $\text{Fr}(U)$  denotes the frontier of  $U$ .

Examples 2.17 and 2.18 show the independence of complementary weakly  $eR$ -openness and weakly  $eR$ -openness.

**Example 2.17** Let  $X = \{a, b, c, d\}$  and

$$\tau = \{\emptyset, X, \{a, d\}\} \quad \text{and} \quad \sigma = \{\emptyset, \{a\}, \{b\}, \{a, b\}, \{a, c\}, \{a, b, c\}, \{a, b, d\}, X\}.$$

The identity function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is weakly  $eR$ -open, but it is not c.w. $eR$ -o.

**Example 2.18** Let  $X = \{a, b\}$ ,  $\tau = \{\emptyset, X, \{a\}, \{b\}\}$  and  $\sigma = \{\emptyset, X, \{b\}\}$ . The identity function  $f : (X, \tau) \rightarrow (X, \sigma)$  is c.w. $eR$ -o., but it is not weakly  $eR$ -open.

**Theorem 2.19** If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is bijective weakly  $eR$ -open and c.w. $eR$ -o, then  $f$  is  $e$ - $\theta$ -open.

**Proof.** Let  $U$  be an open subset in  $X$  with  $x \in U$ . Since  $f$  is weakly  $eR$ -open, by Theorem 2.7(i) there exists an  $e$ - $\theta$ -open set  $V$  containing  $f(x) = y$  such that  $V \subset f(cl(U))$ . Now  $Fr(U) = cl(U) \setminus U$  and thus  $x \notin Fr(U)$ . Hence  $y \notin f(Fr(U))$  and therefore  $y \in V \setminus f(Fr(U))$ . Put  $V_y = V \setminus f(Fr(U))$ . Now  $V_y$  is an  $e$ - $\theta$ -open set since  $f$  is c.w. $eR$ -o. Since  $y \in V_y$ , then  $y \in f(cl(U))$ . But  $y \notin f(Fr(U))$  and thus  $y \in f(cl(U)) \setminus f(Fr(U)) = f(U)$  which implies that  $y \in f(U)$ . Therefore  $f(U) = \cup\{V_y | (V_y \in e\theta O(Y))(y \in f(U))\}$ . Hence  $f$  is  $e$ - $\theta$ -open. ■

Recall that a space  $X$  is said to be  $e$ -connected [3] if  $X$  is not the union of two disjoint nonempty  $e$ -open sets.

**Theorem 2.20** If  $f : (X, \tau) \rightarrow (Y, \sigma)$  is a bijective weakly  $eR$ -open of a space  $X$  onto an  $e$ -connected space  $Y$ , then  $X$  is connected.

**Proof.** Let  $f$  be a bijective weakly  $eR$ -open of a space  $X$  onto an  $e$ -connected space  $Y$  and suppose that  $X$  is not connected.

$$\begin{aligned} X \text{ is not connected} &\Rightarrow (\exists U_1, U_2 \in \tau \setminus \{\emptyset\})(U_1 \cap U_2 = \emptyset)(U_1 \cup U_2 = X) \left. \vphantom{\exists U_1, U_2} \right\} \Rightarrow \\ &\quad f \text{ is bijective weakly } eR\text{-open} \\ &\Rightarrow (f(U_i) \in \sigma \setminus \{\emptyset\})(\cap_i f(U_i) = \emptyset)(\cup_i f(U_i) = Y)(f(U_i) \subset e\text{-int}_\theta(f(cl(U_i))) = e\text{-int}_\theta(f(U_i))) \quad (i = 1, 2) \\ &\Rightarrow (f(U_i) \in \sigma \setminus \{\emptyset\})(\cap_i f(U_i) = \emptyset)(\cup_i f(U_i) = Y)(f(U_i) = e\text{-int}_\theta(f(U_i))) \quad (i = 1, 2) \\ &\Rightarrow (f(U_i) \in e\theta O(Y) \setminus \{\emptyset\})(\cap_i f(U_i) = \emptyset)(\cup_i f(U_i) = Y) \quad (i = 1, 2) \end{aligned}$$

Then  $Y$  is not  $e$ -connected which is a contradiction. ■

**Definition 2.21** A space  $X$  is said to be hyperconnected [9] if every nonempty open subset of  $X$  is dense in  $X$ .

**Theorem 2.22** If  $X$  is a hyperconnected space, then a function  $f : (X, \tau) \rightarrow (Y, \sigma)$  is weakly  $eR$ -open if and only if  $f(X)$  is  $e$ - $\theta$ -open in  $Y$ .

**Proof.** *Sufficiency:* Obvious.

*Necessity:* Let  $U$  be a nonempty open subset of  $X$ .

$$\begin{aligned} (U \in \tau)(X \text{ is hyperconnected}) &\Rightarrow cl(U) = X \Rightarrow e\text{-int}_\theta(f(cl(U))) = e\text{-int}_\theta(f(X)) \left. \vphantom{\Rightarrow} \right\} \Rightarrow \\ &\quad f \text{ is weakly } eR\text{-open} \\ &\Rightarrow f(U) \subset f(X) = e\text{-int}_\theta(f(X)) = e\text{-int}_\theta(f(cl(U))). \end{aligned}$$

**Theorem 2.23** Let  $f : (X, \tau) \rightarrow (Y, \sigma)$  be a bijective weakly  $eR$ -open function. Then the following properties hold:

- (a) If  $F$  is  $\theta$ -closed in  $X$ , then  $f(F)$  is  $e$ - $\theta$ -closed in  $Y$ .
- (b) If  $F$  is  $\theta$ -open in  $X$ , then  $f(F)$  is  $e$ - $\theta$ -open in  $Y$ .

**Proof.** (a) Let  $F \in \theta C(X)$ .

$$\begin{aligned} F \in \theta C(X) &\Rightarrow F = cl_\theta(F) \left. \vphantom{\Rightarrow} \right\} \Rightarrow e\text{-cl}_\theta(f(F)) \subset f(cl_\theta(F)) = f(F) \left. \vphantom{\Rightarrow} \right\} \Rightarrow \\ &\quad \text{Theorem 2.8(i)} \quad f(F) \subset e\text{-cl}_\theta(f(F)) \\ &\Rightarrow f(F) = e\text{-cl}_\theta(f(F)) \\ &\Rightarrow f(F) \in e\theta C(Y). \end{aligned}$$

(b) Similarly proved. ■

## Acknowledgements

The authors are grateful to the referee for his/her careful reading and useful comments for the improvement of this paper. This study is dedicated to Professor Dr. Zekeriya



Güney on the occasion of his 67th birthday.

## References

- [1] D.Andrijevic, On  $b$ -open sets. Mat. Vesnik., 48 (1996), 59-64.
- [2] M. Caldas, E. Ekici, S. Jafari, R.M. Latif, On weakly  $BR$ -open functions and their characterizations in topological spaces. Demonstratio Math., 44 (1) (2011), 159-168.
- [3] E. Ekici, New forms of contra continuity. Carpathian J. Math., 24 (1) (2008), 37-45.
- [4] ———, On  $\epsilon$ -open sets,  $\mathcal{DP}^*$ -sets and  $\mathcal{DP}\mathcal{E}^*$ -sets and decompositions of continuity. Arab. J. Sci. Eng. Sect. A Sci., 33 (2) (2008), 269-282.
- [5] N. Levine, Semi-open sets and semi-continuity in topological spaces. Amer. Math. Monthly., 70 (1963), 36-41.
- [6] ———, Strong continuity in topological spaces. Amer. Math. Monthly., 67 (1960), 269.
- [7] A.S. Mashhour, M.E. Abd El-Monsef and S.N. El-Deeb, On precontinuous and weak precontinuous mappings. Proc. Math. Phys. Soc. Egypt., 53 (1982), 47-53.
- [8] O. Njastad, On some classes of nearly open sets. Pacific J. Math., 15 (1965), 961-970.
- [9] T. Noiri, A generalization of closed mappings. Atti. Accad. Naz. Lince Rend. Cl. Sci. Fis. Mat. Natur., 8 (1973), 210-214.
- [10] M. Özkoç and G. Aslm, On strongly  $\theta$ - $\epsilon$ -continuous functions. Bull. Korean Math. Soc., 47 (5) (2010), 1025-1036.
- [11] J.H. Park, Strongly  $\theta$ - $b$ -continuous functions. Acta Math. Hungar., 110 (4) (2006), 347-359.
- [12] M. Stone, Application of the theory of Boolean ring to general topology. Trans. Amer. Math. Soc., 41 (1937), 374-481.
- [13] N.V. Velicko,  $H$ -closed topological spaces. Amer. Math. Soc. Transl., 78 (1968), 103-118.