

On some Frobenius groups with the same prime graph as the almost simple group $\text{PGL}(2, 49)$

A. Mahmoudifar*

^aDepartment of Mathematics, Tehran North Branch, Islamic Azad University, Tehran, Iran.

Received 25 December 2016; Revised 5 March 2017; Accepted 15 April 2017.

Abstract. The prime graph of a finite group G is denoted by $\Gamma(G)$ whose vertex set is $\pi(G)$ and two distinct primes p and q are adjacent in $\Gamma(G)$, whenever G contains an element with order pq . We say that G is unrecognizable by prime graph if there is a finite group H with $\Gamma(H) = \Gamma(G)$, in while $H \not\cong G$. In this paper, we consider finite groups with the same prime graph as the almost simple group $\text{PGL}(2, 49)$. Moreover, we construct some Frobenius groups whose prime graphs coincide with $\Gamma(\text{PGL}(2, 49))$, in particular, we get that $\text{PGL}(2, 49)$ is unrecognizable by its prime graph.

© 2017 IAUCTB. All rights reserved.

Keywords: Almost simple group, prime graph, Frobenius group, element order.

2010 AMS Subject Classification: 20D05, 20D60, 20D08.

1. Introduction

Let N denotes the set of natural numbers. If $n \in N$, then we denote by $\pi(n)$, the set of all prime divisors of n . Let G be a finite group. The set $\pi(|G|)$ is denoted by $\pi(G)$. Also the set of element orders of G is denoted by $\pi_e(G)$. We denote by $\mu(G)$, the maximal numbers of $\pi_e(G)$ under the divisibility relation. The *prime graph* of G is a graph whose vertex set is $\pi(G)$ and two distinct primes p and q are joined by an edge (and we write $p \sim q$), whenever G contains an element of order pq . The prime graph of G is denoted by $\Gamma(G)$. A finite group G is called *unrecognizable by prime graph* if there exists a finite group H such that $\Gamma(H) = \Gamma(G)$, however $H \not\cong G$.

In [10], it is proved that if p is a prime number which is not a Mersenne or Fermat prime and $p \neq 11, 19$ and $\Gamma(G) = \Gamma(\text{PGL}(2, p))$, then G has a unique nonabelian composition

*Corresponding author.
E-mail address: alimahmoudifar@gmail.com.

factor which is isomorphic to $\text{PSL}(2, p)$ and if $p = 13$, then G has a unique nonabelian composition factor which is isomorphic to $\text{PSL}(2, 13)$ or $\text{PSL}(2, 27)$. In [3], it is proved that if $q = p^\alpha$, where p is a prime and $\alpha > 1$ is an integer, then $\text{PGL}(2, q)$ is uniquely determined by its element orders. Also, in [1], it is proved that if $q = p^\alpha$, where p is an odd prime and α is an odd natural number, then $\text{PGL}(2, q)$ is uniquely determined by its prime graph. However, in this paper as the main result we prove that the almost simple group $\text{PGL}(2, 49)$ is unrecognizable by prime graph. Finally we put a question about the existence of Frobenius groups with the same prime graph as the almost simple groups $\text{PGL}(2, q)$.

We note that in this paper, if G is a group, then $\text{Fit}(G)$ is the Fitting subgroup of G that is the product of all nilpotent normal subgroups of G .

2. Preliminary Results

Lemma 2.1 ([17, Lemma 1]) Let G be a finite group and $N \trianglelefteq G$ such that G/N is a Frobenius group with kernel F and cyclic complement C . If $(|F|, |N|) = 1$ and F is not contained in $NC_G(N)/N$, then $p|C| \in \pi_e(G)$ for some prime divisor p of $|N|$.

Lemma 2.2 ([8, Main Theorem]) Let G be a finite group and $|\pi(G)| \geq 3$. If there exist prime numbers $r, s, t \in \pi(G)$, such that $\{tr, ts, rs\} \cap \pi_e(G) = \emptyset$, then G is non-solvable.

We note that a Z -group is a group in which every sylow subgroup is cyclic. Now we have the following lemma:

Lemma 2.3 ([19, Theorem 18.6]) Let G be a nonsolvable Frobenius complement. Then G has a normal subgroup G_0 with $|G : G_0| = 1$ or 2 such that $G_0 = \text{SL}(2, 5) \times M$ with M a Z -group of order prime to $2, 3$ and 5 .

A finite group G is called a 2-Frobenius group when there exists a normal series, $1 \leq H \leq K \leq G$, such that K and G/H are Frobenius groups with kernels H and K/H , respectively. Using [14, Theorem A], we have the following result:

Lemma 2.4 Let G be a finite group with $t(G) \geq 2$. Then one of the following holds:

- (a) G is a Frobenius or 2-Frobenius group;
- (b) there exists a nonabelian simple group S such that $S \leq \overline{G} := G/N \leq \text{Aut}(S)$ for some nilpotent normal subgroup N of G .

Lemma 2.5 ([20]) Let $G = L_n^\epsilon(q)$, $q = p^m$, be a simple group which acts absolutely irreducibly on a vector space W over a field of characteristic p . Denote $H = W \rtimes G$. If $n = 2$ and q is odd then $2p \in \pi_e(H)$.

3. Main Results

Lemma 3.1 There are infinitely many finite Frobenius group G such that $\Gamma(G) = \Gamma(\text{PGL}(2, 49))$.

Proof. Let F be a finite field of characteristic 7. Also there are some elements α and β included in F such that $\alpha^2 = -1$ and $\beta^2 = 5$. We know that such a finite field does exist and moreover there are infinitely many field with these properties.

Now we construct some linear groups as follow:

$$C := \left\langle \begin{pmatrix} -1 & 1 & 0 \\ -1 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & \alpha & 0 \\ \alpha & \frac{\beta+1}{2} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right\rangle,$$

$$K := \left\langle \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 1 \end{pmatrix} \right\rangle.$$

By the above definition, $C \cong \langle x, y, z | x^3 = y^5 = z^2 = 1, x^z = z, y^z = y, (xy)^2 = z \rangle$. This implies that $C \cong \text{SL}(2, 5)$. Also we have $K \cong F \oplus F$, is a direct sum of additive group F by itself. This means K is isomorphic to a vector space of dimension 2 over F and so $|K| = |F|^2$. It is obvious that C belongs to the normalizer of K in $\text{GL}(3, F)$.

Now we define $G := K \rtimes C$. Since K is an elementary abelian 7-group, it is easy to prove that C acts fixed point freely on K by conjugation. Hence G is a Frobenius group with kernel K and complement C . This implies that in the prime graph of G , 7 is an isolated vertex. Also by $\Gamma(\text{SL}(2, 5))$, we get that 2 is adjacent to 3 and 5 and there is no edge between 3 and 5 in $\Gamma(G)$. Therefore, $\Gamma(G)$ coincides to $\Gamma(\text{PGL}(2, 49))$, which completes the proof. ■

Lemma 3.2 Each following group G is an almost simple group related to the simple group S . Moreover, G has a prime graph which coincide with the prime graph of the almost simple group $\text{PGL}(2, 49)$:

- (1) $G = S_7$ and $S = A_7$.
- (2) $G = U_4(3) \cdot 2$ and $S = U_4(3)$.
- (3) $G = U_3(5)$ or $G = U_3(5) \cdot 2$ and $S = U_3(5)$.

Proof. Using [4], it is straightforward. ■

Theorem 3.3 Let G be a finite group with the prime graph as same as the prime graph of $\text{PGL}(2, 49)$. Then G is isomorphic to one of the following groups:

- (1) A Frobenius group $K \rtimes C$, such that K is a 7-group and C contains a subgroup C_0 whose index in C is at most 2 and C_0 is isomorphic to $\text{SL}(2, 5)$.
- (2) One of the almost simple groups: $S_7, U_4(3) \cdot 2, U_3(5) \cdot 2$ or $\text{PGL}(2, 49)$.
- (3) The simple group: $U_3(5)$.

In particular, $\text{PGL}(2, 49)$ is unrecognizable by prime graph.

Proof. By [18, Lemma 7], it follows that $\mu(\text{PGL}(2, 49)) = \{7, 48, 50\}$. Hence, the connected components of the prime graph of $\text{PGL}(2, 49)$ are exactly $\{7\}$ and $\{2, 3, 5\}$. Also by $\mu(\text{PGL}(2, 49))$, there is no edge between 3 and 5 in $\Gamma(\text{PGL}(2, 49))$. Now since $\Gamma(G) = \Gamma(\text{PGL}(2, 49))$, we deduce that these relations hold in the prime graph of G .

First we claim that G is not solvable. On the contrary, let G be a solvable group. So there is a Hall $\{3, 5, 7\}$ -subgroup in G , say H . On the other hand $\{3, 5, 7\}$ is an independent subset of $\Gamma(G)$, which is a contradiction by Lemma 2.2. Therefore, G is not solvable and so by Lemma 2.4, either G is a Frobenius group or there is a nonabelian simple group S such that $S \leq \bar{G} := G/\text{Fit}(G) \leq \text{Aut}(S)$.

Let G be a Frobenius group with kernel K and complement C . By Lemma 2.3, we know that K is nilpotent and $\pi(C)$ is a connected component of the prime graph of G . Hence we conclude that $\pi(K) = \{7\}$ and $\pi(C) = \{2, 3, 5\}$, since 7 is an isolated vertex in

$\Gamma(G)$. Hence if C is solvable, then G is a solvable which is a contradiction by the above argument.

Thus we suppose that C is non-solvable. Then by Lemma 2.3, the complement C has a normal subgroup C_0 with index at most 2 which is isomorphic to $\text{SL}(2, 5) \times M$, where $\pi(M) \cap \{2, 3, 5\} = \emptyset$. On the other hand, by the previous argument, we know that $\pi(C) = \{2, 3, 5\}$. This implies that $M = 1$ and so $C_0 \cong \text{SL}(2, 5)$. Also by Lemma 3.1, we know that this such Frobenius complement exists. Hence G can be isomorphic to a Frobenius group $K : C$, where K is a 7-subgroup and C contains a subgroup isomorphic to $\text{SL}(2, 5)$ whose index is at most 2, Therefore if G is a Frobenius group, then we get Case (1).

Now we may assume that G is neither a Frobenius nor a 2-Frobenius group. Hence by Lemma 2.4, there exists a nonabelian simple group S such that:

$$S \leq \bar{G} := G/K \leq \text{Aut}(S)$$

in which K is the Fitting subgroup of G . Since $\{2, 7\}$ is an independent subset of $\Gamma(G)$, by Lemma 2.4, we conclude that $7 \in \pi(S)$ and $7 \notin \pi(K) \cup \pi(\bar{G}/S)$. Also we know that $\pi(S) \subseteq \pi(G)$. Since $\pi(G) = \{2, 3, 5, 7\}$, so by [13, Table 8], we get that S is isomorphic to $A_7, A_8, A_9, A_{10}, S_6(2), O_8^+(2), L_3(2^2), L_2(2^3), U_3(3), U_4(3), U_3(5), L_2(7), S_4(7), L_2(7^2)$ or J_2 . Now we consider each possibility for the simple group S .

Let $S \cong L_2(7)$. Then $5 \in \pi(K)$, since $5 \notin (\pi(S) \cup \pi(\bar{G}/S))$. On the other hand S contains a $\{3, 7\}$ -subgroup H . Hence G has a subgroup isomorphic to $K_5 : H$ where K_5 is 5-group. On the other hand $K_5 : H$ is solvable and so there is an edge between to prime numbers in $\Gamma(K_5 : H)$, which is impossible since $\Gamma(K_5 : H)$ is a subgraph of $\Gamma(G)$. Thus $S \not\cong L_2(7)$.

Let $S \cong L_2(2^3)$. In this case, $5 \in \pi(K)$. Also we know that S contains a Frobenius group isomorphic to $8 : 7$. Hence by Lemma 2.1, we get that G has an element order $5 \cdot 7$, which is a contradiction.

Let $S \cong A_8, A_9$ or A_{10} . Thus S consists an element of order $3 \cdot 5$, which contradicts to the prime graph of G .

Let $S \cong J_2, O_8^+(2)$ or $S_6(2)$. In this case S contains an element of order 15, which is a contradiction.

By Lemma 3.2, the finite group S can be isomorphic to each simple group $A_7, U_3(3), U_4(3)$ and $U_3(5)$.

Let S be isomorphic to $\text{PSL}_2(49)$. Hence $\text{PSL}_2(49) \leq \bar{G} \leq \text{Aut}(\text{PSL}_2(49))$.

Let $\pi(K)$ contains a prime r such that $r \neq 7$. Since K is nilpotent, we may assume that K is a vector space over a field with r elements (analogous to the proof of Lemma 3.1). Hence the prime graph of the semidirect product $K \rtimes \text{PSL}_2(49)$ is a subgraph of $\Gamma(G)$. Let B be a Sylow 7-subgroup of $\text{PSL}_2(49)$. We know that B is not cyclic. On the other hand $K \rtimes B$ is a Frobenius group such that $\pi(K \rtimes B) = \{r, 7\}$. Hence B should be cyclic which is a contradiction. This implies that $K = 1$, since $7 \notin \pi(K)$.

We know that $\text{Out}(\text{PSL}_2(49)) \cong Z_2 \times Z_2$. Since in the prime graph of $\text{PSL}_2(49)$ there is not any edge between 2 and 5, we get that $G \not\cong \text{PSL}_2(49)$. If $G = \text{PSL}_2(49) : \langle \theta \rangle$, where θ is a field automorphism, then we get that 2 and 7 are adjacent in G (for example see Page 26 of [1]), which is a contradiction. Also if $G = \text{PSL}_2(49) : \langle \gamma \rangle$, where γ is a diagonal automorphism, then we get that G is isomorphic to $\text{PGL}_2(49)$, which completes the proof. ■

Problem. Let $G = \text{PGL}(2, q)$ be an almost simple group related to the simple group $\text{PSL}(2, q)$. Find all Frobenius group H such that $\Gamma(H) = \Gamma(G)$.

Acknowledgements

The financial support devoted to this project from Islamic Azad University (Tehran North Branch) is deeply appreciated by the author. In addition, the author would like to give his sincere thanks to the referee for being so precise in making valuable suggestions.

References

- [1] Z. Akhlaghi, M. Khatami, B. Khosravi, Characterization by prime graph of $PGL(2, p^k)$ where p and k are odd, International Journal of Algebra and Computation. 20 (2010), 847-873.
- [2] A. A. Buturlakin, Spectra of Finite Symplectic and Orthogonal Groups, Siberian Advances in Mathematics. 21 (2011), 176-210.
- [3] G. Y. Chen, V. D. Mazurov, W. J. Shi, A. V. Vasil'ev, A. Kh. Zhurtov, Recognition of the finite almost simple groups $PGL_2(q)$ by their spectrum, J. Group Theory. 10 (2007), 71-85.
- [4] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, R. A. Wilson, Atlas of Finite Groups, Oxford University Press, Oxford, 1985.
- [5] M.A. Grechkoseva, On element orders in covers of finite simple classical groups, J. Algebra. 339 (2011), 304-319.
- [6] D. Gorenstein, Finite Groups, Harper and Row, New York, 1968.
- [7] M. Hagie, The prime graph of a sporadic simple group, Comm. Algebra. 31 (2003), 4405-4424.
- [8] G. Higman, Finite groups in which every element has prime power order, J. London Math. Soc. 32 (1957), 335-342.
- [9] M. Khatami, B. Khosravi, Z. Akhlaghi, NCF-distinguishability by prime graph of $PGL(2, p)$, where p is a prime, Rocky Mountain J. Math. 41 (2011), 1523-1545.
- [10] B. Khosravi, n -Recognition by prime graph of the simple group $PSL(2, q)$, J. Algebra. Appl. 7 (2008), 735-748.
- [11] B. Khosravi, 2-Recognizability of $PSL(2, p^2)$ by the prime graph, Siberian Math. J. 49 (2008), 749-757.
- [12] B. Khosravi, B. Khosravi, B. Khosravi, On the prime graph of $PSL(2, p)$ where $p > 3$ is a prime number, Acta. Math. Hungarica. 116 (2007), 295-307.
- [13] R. Kogani-Moghadam, A. R. Moghaddamfar, Groups with the same order and degree pattern, Sci. China. Math. 55 (2012), 701-720.
- [14] A. S. Kondrat'ev, Prime graph components of finite simple groups, Math. USSR-SB. 67 (1990), 235-247.
- [15] A. Mahmoudifar, B. Khosravi, On quasirecognition by prime graph of the simple groups $A_n^+(p)$ and $A_n^-(p)$, J. Algebra. Appl. 14 (2015), 12 pages.
- [16] A. Mahmoudifar, B. Khosravi, On the characterization of alternating groups by order and prime graph, Sib. Math. J. 56 (2015), 125-131.
- [17] V. D. Mazurov, Characterizations of finite groups by sets of their element orders, Algebra Logic. 36 (1997), 23-32.
- [18] A. R. Moghaddamfar, W. J. Shi, The number of finite groups whose element orders is given, Beitrage Algebra Geom. 47 (2006), 463-479.
- [19] D. S. Passman, Permutation groups, W. A. Benjamin, New York, 1968.
- [20] A. V. Zavarnitsine, Fixed points of large prime-order elements in the equicharacteristic action of linear and unitary groups, Sib. Electron. Math. Rep. 8 (2011), 333-340.