

Interval valued fuzzy weak bi-ideals of Γ -near-rings

V. Chinnadurai^{a,*}, K. Arulmozhi^a, S. Kadarasi^a

^aDepartment of Mathematics, Annamalai University, Annamalainagar-608 002, India.

Received 11 May 2017; Revised 25 September 2017; Accepted 1 November 2017.

Abstract. In this paper, we introduce the concept of interval valued fuzzy weak bi-ideals of Γ -near-rings, which is a generalized concept of fuzzy weak bi-ideals of Γ -near-rings. We also characterize some properties and provide examples of interval valued fuzzy weak bi-ideals of Γ -near-rings.

© 2017 IAUCTB. All rights reserved.

Keywords: Γ -near-rings, fuzzy weak bi-ideals, interval valued fuzzy weak bi-ideals, homomorphism and anti-homomorphism.

2010 AMS Subject Classification: 16Y30, 03E72, 08A72.

1. Introduction

Zadeh introduced the concept of fuzzy sets in 1965 [21], and also, generalized it to interval valued fuzzy subsets [22]. Near-ring was introduced by Pilz [14]. Gamma -near-ring was introduced by Satyanarayana [15] in 1984. The concept of bi-ideals of near-ring was applied to Γ -near-rings by Tamizh chelvam et al. [16]. The idea of fuzzy ideals of near-rings was proposed by Kim et al. [9]. Fuzzy ideals in *Gamma*-near-rings proposed by Jun et al. [8] in 1998. Moreover, Manikantan [10] introduced the notion of fuzzy bi-ideals of near-rings and discussed some properties. Meenakumari et al. [12] studied the fuzzy bi-ideals in gamma -near-rings. Cho et al. [20] introduced the concept of weak bi-ideals of near-rings. Thillaigovindan *at al.* [17] studied the interval valued fuzzy quasi-ideals of semigroups. Chinnadurai et al. [3] studied the fuzzy weak bi-ideals of near rings. Thillaigovindan et al. [18] worked on interval valued fuzzy ideals of near-rings.

*Corresponding author.

E-mail address: kv.chinnadurai@yahoo.com (V. Chinnadurai); arulmozhiems@gmail.com (K. Arulmozhi); kadalarsi89@gmail.com (S. Kadarasi).

In this paper, we define a new notion of an interval valued fuzzy weak bi-ideals of Γ -near-rings, which is a generalized concept of interval valued fuzzy bi-ideals of Γ -near-rings. We also investigate some of its properties and illustrate with examples.

2. Preliminaries

In this section, we list some basic definitions.

Definition 2.1 [14] A near-ring is an algebraic system $(R, +, \cdot)$ consisting of a non empty set R together with two binary operations called $+$ and \cdot such that $(R, +)$ is a group not necessarily abelian and (R, \cdot) is a semigroup connected by the following distributive law: $(x + z) \cdot y = x \cdot y + z \cdot y$ valid for all $x, y, z \in R$. We use the word ‘near-ring’ to mean ‘right near-ring’. We denote xy instead of $x \cdot y$.

Definition 2.2 [15] A Γ -near-ring is a triple $(M, +, \Gamma)$ where

- (i) $(M, +)$ is a group,
- (ii) Γ is a nonempty set of binary operators on M such that for each $\alpha \in \Gamma$, $(M, +, \alpha)$ is a near-ring,
- (iii) $x\alpha(y\beta z) = (x\alpha y)\beta z$ for all $x, y, z \in M$ and $\alpha, \beta \in \Gamma$.

Definition 2.3 [12] A Γ -near-ring M is said to be zero-symmetric if $x\alpha 0 = 0$ for all $x \in M$ and $\alpha \in \Gamma$.

Throughout this paper M denotes a zero-symmetric right Γ -near-ring with atleast two elements.

Definition 2.4 [15] A subset A of a Γ -near-ring M is called a left (resp. right) ideal of M if

- (i) $(A, +)$ is a normal subgroup of $(M, +)$, (i.e) $x - y \in A$ for all $x, y \in A$ and $y + x - y \in A$ for $x \in A, y \in M$
- (ii) $u\alpha(x + v) - u\alpha v \in A$ (resp. $x\alpha u \in A$) for all $x \in A, \alpha \in \Gamma$ and $u, v \in M$.

Definition 2.5 [15] Let M be a Γ -near-ring. Given two subsets A and B of M , we define $A\Gamma B = \{a\alpha b \mid a \in A, b \in B \text{ and } \alpha \in \Gamma\}$ and also define another operation $*$ on the class of subset of M define by $A\Gamma * B = \{a\gamma(a' + b) - a\gamma a' \mid a, a' \in A, \gamma \in \Gamma, b \in B\}$.

Definition 2.6 [16] A subgroup B of $(M, +)$ is called a bi-ideal of M if and only if $B\Gamma M\Gamma B \subseteq B$.

Definition 2.7 [5] A subgroup H of $(M, +)$ is said to be a weak bi-ideal of M if $H\Gamma H\Gamma H \subseteq H$.

The characteristic function of M is denoted by \mathbf{M} .

Definition 2.8 [22] If X be any set. A mapping $\eta : X \rightarrow D[0, 1]$ is called an interval valued fuzzy subset (briefly, an i.v fuzzy subset) of X , where $D[0, 1]$ denotes the family of closed subintervals of $[0, 1]$ and $\tilde{\eta}(x) = [\eta^-(x), \eta^+(x)]$ for all $x \in X$, where $\eta^-(x)$ and $\eta^+(x)$ are fuzzy subsets of X such that $\eta^-(x) \leq \eta^+(x)$ for all $x \in X$.

Definition 2.9 [17] By an interval number \tilde{a} , we mean an interval $[a^-, a^+]$ such that $0 \leq a^- \leq a^+ \leq 1$ and where a^- and a^+ are the lower and upper limits of \tilde{a} respectively. The set of all closed subintervals of $[0, 1]$ is denoted by $D[0, 1]$. We also identify the interval $[a, a]$ by the number $a \in [0, 1]$. For any interval numbers $\tilde{a}_j = [a_j^-, a_j^+], \tilde{b}_j = [b_j^-, b_j^+] \in D[0, 1], j \in \Omega$ we define

$\max^i\{\tilde{a}_j, \tilde{b}_j\} = [\max\{a_j^-, b_j^-\}, \max\{a_j^+, b_j^+\}]$, $\min^i\{\tilde{a}_j, \tilde{b}_j\} = [\min\{a_j^-, b_j^-\}, \max\{a_j^+, b_j^+\}]$,
 $\inf^i \tilde{a}_j = [\cap_{j \in \Omega} a_j^-, \cap_{j \in \Omega} a_j^+]$, $\sup^i \tilde{a}_j = [\cup_{j \in \Omega} a_j^-, \cup_{j \in \Omega} a_j^+]$, and let

- (i) $\tilde{a} \leq \tilde{b} \Leftrightarrow a^- \leq b^-$ and $a^+ \leq b^+$,
- (ii) $\tilde{a} = \tilde{b} \Leftrightarrow a^- = b^-$ and $a^+ = b^+$,
- (iii) $\tilde{a} < \tilde{b} \Leftrightarrow \tilde{a} \leq \tilde{b}$ and $\tilde{a} \neq \tilde{b}$,
- (iv) $k\tilde{a} = [ka^-, ka^+]$, whenever $0 \leq k \leq 1$.

Definition 2.10 [17] Let $\tilde{\eta}$ be an i.v fuzzy subset of X and $[t_1, t_2] \in D[0, 1]$. Then the set $\tilde{U}(\tilde{\eta} : [t_1, t_2]) = \{x \in X | \tilde{\eta}(x) \geq [t_1, t_2]\}$ is called the upper level subset of $\tilde{\eta}$.

Definition 2.11 [7, 13, 19] If $\tilde{\eta}$ and $\tilde{\lambda}$ are i.v fuzzy subsets of M . Then $\tilde{\eta} \cap \tilde{\lambda}$, $\tilde{\eta} \cup \tilde{\lambda}$, $\tilde{\eta} + \tilde{\lambda}$, and $\tilde{\eta} * \tilde{\lambda}$ are fuzzy subsets of M defined by:

$$\begin{aligned}
 (\tilde{\eta} \cap \tilde{\lambda})(x) &= \min^i\{\tilde{\eta}(x), \tilde{\lambda}(x)\}. \\
 (\tilde{\eta} \cup \tilde{\lambda})(x) &= \max^i\{\tilde{\eta}(x), \tilde{\lambda}(x)\}. \\
 (\tilde{\eta} + \tilde{\lambda})(x) &= \begin{cases} \sup_{x=y+z}^i\{\min^i\{\tilde{\eta}(y), \tilde{\lambda}(z)\}\} & \text{if } x \text{ is expressible as } x = y + z \\ 0 & \text{otherwise.} \end{cases} \\
 (\tilde{\eta} * \tilde{\lambda})(x) &= \begin{cases} \sup_{x=y\alpha z}^i\{\min^i\{\tilde{\eta}(y), \tilde{\lambda}(z)\}\} & \text{if } x \text{ is expressible as } x = y\alpha z \\ 0 & \text{otherwise.} \end{cases}
 \end{aligned}$$

for $x, y, z \in M$.

Definition 2.12 [5] An i.v fuzzy subset $\tilde{\eta}$ in a Γ -near-ring M is called an i.v fuzzy left (resp. right) ideal of M if

- (i) $\tilde{\eta}$ is an i.v fuzzy normal divisor with respect to the addition,
- (ii) $\tilde{\eta}(u\alpha(x + v) - u\alpha v) \geq \tilde{\eta}(x)$, (resp. $\tilde{\eta}(x\alpha u) \geq \tilde{\eta}(x)$ for all $x, u, v \in M$ and $\alpha \in \Gamma$.

The condition (i) of definition 2.12 means that $\tilde{\eta}$ satisfies:

- (i) $\tilde{\eta}(x - y) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$,
- (ii) $\tilde{\eta}(y + x - y) \geq \tilde{\eta}(x)$, for all $x, y \in M$

Note that $\tilde{\eta}$ is an i.v fuzzy left (resp. right) ideal of Γ -near-ring M , then $\tilde{\eta}(0) \geq \tilde{\eta}(x)$ for all $x \in M$, where 0 is the zero element of M .

Definition 2.13 [6] An i.v fuzzy subset $\tilde{\eta}$ of M is called an i.v fuzzy bi-ideal of M if

- (i) $\tilde{\eta}(x - y) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$ for all $x, y \in M$
- (ii) $\tilde{\eta}(x\alpha y\beta z) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(z)\}$ for all $x, y, z \in M$ and $\alpha, \beta \in \Gamma$.

3. Interval valued fuzzy weak bi-ideals of Γ -near-rings

In this section, we introduce the notion of i.v fuzzy weak bi-ideal of M and discuss some of its properties.

Definition 3.1 An i.v fuzzy set $\tilde{\eta}$ of M is called an i.v fuzzy weak bi-ideal of M , if

- (i) $\tilde{\eta}(x - y) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$ for all $x, y \in M$
- (ii) $\tilde{\eta}(x\alpha y\beta z) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\}$ for all $x, y, z \in M$ and $\alpha, \beta \in \Gamma$.

Example 3.2 Let $M = \{0, a, b, c\}$ be a non-empty set with binary operation $+$ and $\Gamma = \{\alpha, \beta\}$ be a non-empty set of binary operations as shown in the following tables:

+	0	a	b	c
0	0	a	b	c
a	a	0	c	b
b	b	c	0	a
c	c	b	a	0

α	0	a	b	c
0	0	0	0	0
a	a	a	a	a
b	0	0	b	b
c	a	a	c	c

β	0	a	b	c
0	0	0	0	0
a	0	0	0	0
b	0	a	c	b
c	0	a	b	c

Let $\tilde{\eta} : M \rightarrow D[0, 1]$ be an i.v fuzzy subset defined by $\tilde{\eta}(0) = [0.8, 0.9]$, $\tilde{\eta}(a) = [0.6, 0.7]$, $\tilde{\eta}(b) = \tilde{\eta}(c) = [0.2, 0.3]$. Then $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M .

Theorem 3.3 Let $\tilde{\eta}$ be an i.v fuzzy subgroup of M . Then $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M if and only if $\tilde{\eta} * \tilde{\eta} * \tilde{\eta} \subseteq \tilde{\eta}$.

Proof. Assume that $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M . Let $x, y, z, y_1, y_2 \in M$ and $\alpha, \beta \in \Gamma$ such that $x = y\alpha z$ and $y = y_1\beta y_2$. Then

$$\begin{aligned}
(\tilde{\eta} * \tilde{\eta} * \tilde{\eta})(x) &= \sup_{x=y\alpha z}^i \{ \min^i \{ (\tilde{\eta} * \tilde{\eta})(y), \tilde{\eta}(z) \} \} \\
&= \sup_{x=y\alpha z}^i \{ \min^i \{ \sup_{y=y_1\beta y_2}^i \min^i \{ \tilde{\eta}(y_1), \tilde{\eta}(y_2) \}, \tilde{\eta}(z) \} \} \\
&= \sup_{x=y\alpha z}^i \sup_{y=y_1\beta y_2}^i \{ \min^i \{ \min^i \{ \tilde{\eta}(y_1), \tilde{\eta}(y_2) \}, \tilde{\eta}(z) \} \} \\
&= \sup_{x=y_1\beta y_2\alpha z}^i \{ \min^i \{ \tilde{\eta}(y_1), \tilde{\eta}(y_2), \tilde{\eta}(z) \} \} \\
&\quad (\text{since } \tilde{\eta} \text{ is an i.v fuzzy weak bi-ideal of } M, \tilde{\eta}(y_1\beta y_2\alpha z) \geq \min^i \{ \tilde{\eta}(y_1), \tilde{\eta}(y_2), \tilde{\eta}(z) \}) \\
&\leq \sup_{x=y_1\beta y_2\alpha z}^i \tilde{\eta}(y_1\beta y_2\alpha z) \\
&= \tilde{\eta}(x).
\end{aligned}$$

If x can not be expressed as $x = y\alpha z$, then $(\tilde{\eta} * \tilde{\eta} * \tilde{\eta})(x) = \tilde{0} \leq \tilde{\eta}(x)$. In both cases $\tilde{\eta} * \tilde{\eta} * \tilde{\eta} \subseteq \tilde{\eta}$. Conversely, assume that $\tilde{\eta} * \tilde{\eta} * \tilde{\eta} \subseteq \tilde{\eta}$. For $x', x, y, z \in M$ and $\alpha, \beta, \alpha_1, \beta_1 \in \Gamma$. Let x' be such that $x' = x\alpha y\beta z$. Then

$$\begin{aligned}
\tilde{\eta}(x\alpha y\beta z) &= \eta(x') \geq (\tilde{\eta} * \tilde{\eta} * \tilde{\eta})(x') \\
&= \sup_{x'=p\alpha_1 q}^i \{ \min^i \{ (\tilde{\eta} * \tilde{\eta})(p), \tilde{\eta}(q) \} \} \\
&= \sup_{x'=p\alpha_1 q}^i \{ \min^i \{ \sup_{p=p_1\beta_1 p_2}^i \min^i \{ \tilde{\eta}(p_1), \tilde{\eta}(p_2) \}, \tilde{\eta}(q) \} \} \\
&= \sup_{x'=p_1\beta_1 p_2\alpha_1 q}^i \{ \min^i \{ \tilde{\eta}(p_1), \tilde{\eta}(p_2), \tilde{\eta}(q) \} \} \\
&\geq \min^i \{ \tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z) \}.
\end{aligned}$$

Hence $\tilde{\eta}(x\alpha y\beta z) \geq \min^i \{ \tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z) \}$. ■

Lemma 3.4 Let $\tilde{\eta}$ and $\tilde{\lambda}$ be an i.v fuzzy weak bi-ideals of M . Then the products $\tilde{\eta} * \tilde{\lambda}$ and $\tilde{\lambda} * \tilde{\eta}$ are also an i.v fuzzy weak bi-ideals of M .

Proof. Let $\tilde{\eta}$ and $\tilde{\lambda}$ be an i.v fuzzy weak bi-ideals of M and let $\alpha, \alpha_1, \alpha_2 \in \Gamma$. Now

$$\begin{aligned} (\tilde{\eta} * \tilde{\lambda})(x - y) &= \sup_{x-y=a\alpha b}^i \min^i \{ \tilde{\eta}(a), \tilde{\lambda}(b) \} \\ &\geq \sup_{x-y=a_1\alpha_1 b_1 - a_2\alpha_2 b_2 < (a_1 - a_2)(b_1 - b_2)}^i \min^i \{ \tilde{\eta}(a_1 - a_2), \tilde{\lambda}(b_1 - b_2) \} \\ &\geq \sup^i \min^i \{ \min^i \{ \tilde{\eta}(a_1), \tilde{\eta}(a_2) \}, \min^i \{ \tilde{\lambda}(b_1), \tilde{\lambda}(b_2) \} \} \\ &= \sup^i \min^i \{ \min^i \{ \tilde{\eta}(a_1), \tilde{\lambda}(b_1) \}, \min^i \{ \tilde{\eta}(a_2), \tilde{\lambda}(b_2) \} \} \\ &\geq \min^i \{ \sup_{x=a_1\alpha_1 b_1}^i \min^i \{ \tilde{\eta}(a_1), \tilde{\lambda}(b_1) \}, \sup_{y=a_2\alpha_2 b_2}^i \min^i \{ \tilde{\eta}(a_2), \tilde{\lambda}(b_2) \} \} \\ &= \min^i \{ (\tilde{\eta} * \tilde{\lambda})(x), (\tilde{\eta} * \tilde{\lambda})(y) \}. \end{aligned}$$

It follows that $\tilde{\eta} * \tilde{\lambda}$ is an i.v fuzzy subgroup of M . Further,

$$\begin{aligned} (\tilde{\eta} * \tilde{\lambda}) * (\tilde{\eta} * \tilde{\lambda}) * (\tilde{\eta} * \tilde{\lambda}) &= \tilde{\eta} * \tilde{\lambda} * (\tilde{\eta} * \tilde{\lambda} * \tilde{\eta}) * \tilde{\lambda} \\ &\subseteq \tilde{\eta} * \tilde{\lambda} * (\tilde{\lambda} * \tilde{\lambda} * \tilde{\lambda}) * \tilde{\lambda}, \text{ since } \tilde{\lambda} \text{ is an i.v fuzzy weak bi-ideal of } M \\ &\subseteq \tilde{\eta} * (\tilde{\lambda} * \tilde{\lambda} * \tilde{\lambda}), \text{ since } \tilde{\lambda} \text{ is an i.v fuzzy weak bi-ideal of } M \\ &\subseteq \tilde{\eta} * \tilde{\lambda}. \end{aligned}$$

Therefore $\tilde{\eta} * \tilde{\lambda}$ is an i.v fuzzy weak bi-ideal of M . Similarly $\tilde{\lambda} * \tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M . ■

Lemma 3.5 Every i.v fuzzy ideal of M is an i.v fuzzy bi-ideal of M .

Proof. Let $\tilde{\eta}$ be an i.v fuzzy ideal of M . Then

$$\tilde{\eta} * \mathbf{M} * \tilde{\eta} \subseteq \tilde{\eta} * \mathbf{M} * \mathbf{M} \subseteq \tilde{\eta} * \mathbf{M} \subseteq \tilde{\eta}$$

since $\tilde{\eta}$ is an i.v fuzzy ideal of M . This implies that $\tilde{\eta} * \mathbf{M} * \tilde{\eta} \subseteq \tilde{\eta}$. Therefore $\tilde{\eta}$ is an i.v fuzzy bi-ideal of M . ■

Theorem 3.6 Every i.v fuzzy bi-ideal of M is an i.v fuzzy weak bi-ideal of M .

Proof. Assume that $\tilde{\eta}$ is an i.v fuzzy bi-ideal of M . Then $\tilde{\eta} * \mathbf{M} * \tilde{\eta} \subseteq \tilde{\eta}$. We have $\tilde{\eta} * \tilde{\eta} * \tilde{\eta} \subseteq \tilde{\eta} * \mathbf{M} * \tilde{\eta}$. This implies that $\tilde{\eta} * \tilde{\eta} * \tilde{\eta} \subseteq \tilde{\eta} * \mathbf{M} * \tilde{\eta} \subseteq \tilde{\eta}$. Therefore $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M . ■

Theorem 3.7 Every i.v fuzzy ideal of M is an i.v fuzzy weak bi-ideal of M .

Proof. By Lemma 3.5, every i.v fuzzy ideal of M is an i.v fuzzy bi-ideal of M . By Theorem 3.6, every i.v fuzzy bi-ideal of M is an i.v fuzzy weak bi-ideal of M . Therefore $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M . ■

However the converse of the Theorems 3.6 and 3.7 is not true in general which is demonstrated by the following example.

Example 3.8 Let $M = \{0, a, b, c\}$ be a non-empty set with binary operation $+$ and $\Gamma = \{\alpha, \beta\}$ be a nonempty set of binary operations as shown in the following tables:

+	0	a	b	c
0	0	a	b	c
a	a	0	c	b
b	b	c	0	a
c	c	b	a	0

α	0	a	b	c
0	0	0	0	0
a	0	a	0	a
b	0	0	b	b
c	0	a	b	c

β	0	a	b	c
0	0	0	0	0
a	0	0	0	0
b	0	a	c	b
c	0	a	b	c

Let $\tilde{\eta} : M \rightarrow D[0, 1]$ be an i.v fuzzy subset defined by $\tilde{\eta}(0) = [0.7, 0.8], \tilde{\eta}(a) = [0.3, 0.4] = \tilde{\eta}(b)$ and $\tilde{\eta}(c) = [0.5, 0.6]$. Then $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M . But $\tilde{\eta}$ is not an i.v fuzzy ideal and bi-ideal of M , since $\tilde{\eta}(bac) = \tilde{\eta}(b) = [0.3, 0.4] \not\supseteq [0.5, 0.6] = \tilde{\eta}(c), \tilde{\eta}(b\beta c) = \tilde{\eta}(b) = [0.3, 0.4] \not\supseteq [0.5, 0.6] = \tilde{\eta}(c)$ and $\tilde{\eta}(c\alpha b\beta c) = \tilde{\eta}(b) = [0.3, 0.4] \not\supseteq [0.5, 0.6] = \min^i\{\tilde{\eta}(c), \tilde{\eta}(c)\}$.

Theorem 3.9 Let $\{\tilde{\eta}_i | i \in \Omega\}$ be family of i.v fuzzy weak bi-ideals of a Γ - near-ring M , then $\bigcap_{i \in \Omega} \tilde{\eta}_i$ is also an i.v fuzzy weak bi-ideal of M , where Ω is any index set.

Proof. Let $\{\tilde{\eta}_i | i \in \Omega\}$ be a family of i.v fuzzy weak bi-ideals of M . Let $x, y, z \in M, \alpha, \beta \in \Gamma$ and $\tilde{\eta} = \bigcap_{i \in \Omega} \tilde{\eta}_i$. Then, $\tilde{\eta}(x) = \bigcap_{i \in \Omega} \tilde{\eta}_i(x) = (\inf_{i \in \Omega}^i \tilde{\eta}_i)(x) = \inf_{i \in \Omega}^i \tilde{\eta}_i(x)$. Now,

$$\begin{aligned} \tilde{\eta}(x - y) &= \inf_{i \in \Omega}^i \tilde{\eta}_i(x - y) \\ &\geq \inf_{i \in \Omega}^i \min^i\{\tilde{\eta}_i(x), \tilde{\eta}_i(y)\} \\ &= \min^i\{\inf_{i \in \Omega}^i \tilde{\eta}_i(x), \inf_{i \in \Omega}^i \tilde{\eta}_i(y)\} \\ &= \min^i\left\{\bigcap_{i \in \Omega} \tilde{\eta}_i(x), \bigcap_{i \in \Omega} \tilde{\eta}_i(y)\right\} \\ &= \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}, \end{aligned}$$

and

$$\begin{aligned} \tilde{\eta}(x\alpha y\beta z) &= \inf_{i \in \Omega}^i \tilde{\eta}_i(x\alpha y\beta z) \\ &\geq \inf_{i \in \Omega}^i \min^i\{\tilde{\eta}_i(x), \tilde{\eta}_i(y), \tilde{\eta}_i(z)\} \\ &= \min^i\{\inf_{i \in \Omega}^i \tilde{\eta}_i(x), \inf_{i \in \Omega}^i \tilde{\eta}_i(y), \inf_{i \in \Omega}^i \tilde{\eta}_i(z)\} \\ &= \min^i\left\{\bigcap_{i \in \Omega} \tilde{\eta}_i(x), \bigcap_{i \in \Omega} \tilde{\eta}_i(y), \bigcap_{i \in \Omega} \tilde{\eta}_i(z)\right\} \\ &= \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\}. \end{aligned}$$

■

Theorem 3.10 Let $\tilde{\eta}$ be an i.v fuzzy subset of M . Then $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M if and only if $\tilde{U}(\tilde{\eta} : [t_1, t_2])$ is a weak bi-ideal of M , for all $[t_1, t_2] \in D[0, 1]$.

Proof. Assume that $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M . Let $[t_1, t_2] \in D[0, 1]$ such that $x, y \in \tilde{U}(\tilde{\eta} : [t_1, t_2])$. Then $\tilde{\eta}(x - y) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\} \geq \min^i\{[t_1, t_2], [t_1, t_2]\} = [t_1, t_2]$. Thus $x - y \in \tilde{U}(\tilde{\eta} : [t_1, t_2])$. Let $x, y, z \in \tilde{U}(\tilde{\eta} : [t_1, t_2])$ and $\alpha, \beta \in \Gamma$. We have $\tilde{\eta}(x\alpha y\beta z) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\} \geq \min^i\{[t_1, t_2], [t_1, t_2], [t_1, t_2]\} = [t_1, t_2]$. Therefore $x\alpha y\beta z \in \tilde{U}(\tilde{\eta} : [t_1, t_2])$. Hence $\tilde{U}(\tilde{\eta} : [t_1, t_2])$ is a weak bi-ideal of M .

Conversely, assume $\tilde{U}(\tilde{\eta} : [t_1, t_2])$ is a weak bi-ideal of M , for all $[t_1, t_2] \in D[0, 1]$. Let $x, y \in M$. Suppose $\tilde{\eta}(x - y) < \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$. Choose $[0, 0] < [t_1, t_2] \leq [1, 1]$ such that $\tilde{\eta}(x - y) < [t_1, t_2] < \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$. This implies that $\tilde{\eta}(x) > [t_1, t_2], \tilde{\eta}(y) > [t_1, t_2]$ and $\tilde{\eta}(x - y) < [t_1, t_2]$. Then we have $x, y \in \tilde{U}(\tilde{\eta} : [t_1, t_2])$, but $x - y \notin \tilde{U}(\tilde{\eta} : [t_1, t_2])$ a contradiction. Thus, $\tilde{\eta}(x - y) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$. If there exist $x, y, z \in M$ and $\alpha, \beta \in \Gamma$ such that $\tilde{\eta}(x\alpha y\beta z) < \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\}$. Choose $[t_1, t_2]$ such that $\tilde{\eta}(x\alpha y\beta z) < [t_1, t_2] < \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\}$. Then $\tilde{\eta}(x) > [t_1, t_2], \tilde{\eta}(y) > [t_1, t_2], \tilde{\eta}(z) > [t_1, t_2]$ and $\tilde{\eta}(x\alpha y\beta z) < [t_1, t_2]$. So, $x, y, z \in \tilde{U}(\tilde{\eta} : [t_1, t_2])$, but $x\alpha y\beta z \notin \tilde{U}(\tilde{\eta} : [t_1, t_2])$, which is a contradiction. Hence $\tilde{\eta}(x\alpha y\beta z) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\}$. Therefore, $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M . ■

Theorem 3.11 Let $\tilde{\eta} = [\eta^-, \eta^+]$ be an i.v fuzzy subset of M , then $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of near-ring M if and only if η^-, η^+ are fuzzy weak bi-ideals of M .

Proof. Assume that $\tilde{\eta}$ is an i.v fuzzy weak bi-ideals of near-ring M . For any $x, y, z \in M$ and $\alpha, \beta \in \Gamma$. Now,

$$\begin{aligned} [\eta^-(x - y), \eta^+(x - y)] &= \tilde{\eta}(x - y) \\ &\geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\} \\ &= \min^i\{[\eta^-(x), \eta^+(x)], [\eta^-(y), \eta^+(y)]\} \\ &= \min^i\{[\eta^-(x), \eta^-(y)], \min^i[\eta^+(x), \eta^+(y)]\}. \end{aligned}$$

It follows that $\eta^-(x - y) \geq \min\{\eta^-(x), \eta^-(y)\}$ and $\eta^+(x - y) \geq \min\{\eta^+(x), \eta^+(y)\}$.

$$\begin{aligned} [\eta^-(x\alpha y\beta z), \eta^+(x\alpha y\beta z)] &= \tilde{\eta}(x\alpha y\beta z) \\ &\geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\} \\ &= \min^i\{[\eta^-(x), \eta^+(x)], [\eta^-(y), \eta^+(y)], [\eta^-(z), \eta^+(z)]\} \\ &= \min^i\{[\eta^-(x), \eta^-(y), \eta^-(z)], \min^i[\eta^+(x), \eta^+(y), \eta^+(z)]\}. \end{aligned}$$

It follows that $\eta^-(x\alpha y\beta z) \geq \min\{\eta^-(x), \eta^-(y), \eta^-(z)\}$ and $\eta^+(x\alpha y\beta z) \geq \min\{\eta^+(x), \eta^+(y), \eta^+(z)\}$. Conversely, assume that η^-, η^+ are fuzzy weak bi-ideals of near-ring M . Let $x, y, z \in M$ and $\alpha, \beta \in \Gamma$.

$$\begin{aligned} \eta^-(x - y) &= [\eta^-(x - y), \eta^+(x - y)] \\ &\geq \min\{[\eta^-(x), \eta^-(y)], \min[\eta^+(x), \eta^+(y)]\} \\ &= \min^i\{[\eta^-(x), \eta^+(x)], [\eta^-(y), \eta^+(y)]\} \\ &= \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\} \end{aligned}$$

and

$$\begin{aligned} \tilde{\eta}(x\alpha y\beta z) &= [\eta^-(x\alpha y\beta z), \eta^+(x\alpha y\beta z)] \\ &\geq \min\{[\eta^-(x), \eta^-(y), \eta^-(z)], \min\{\eta^+(x), \eta^+(y), \eta^+(z)\}\} \\ &= \min^i\{[\eta^-(x), \eta^+(x)], [\eta^-(y), \eta^+(y)], [\eta^-(z), \eta^+(z)]\} \\ &= \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\} \end{aligned}$$

Therefore $\tilde{\eta}$ is an i.v fuzzy weak bi-ideals of near-ring M . ■

Theorem 3.12 Let I be a weak bi-ideal of near-ring M then for any $[t_1, t_2] \in D[0, 1]$ with $[t_1, t_2] \neq [0, 0]$, there exists an i.v fuzzy weak bi-ideal $\tilde{\eta}$ of M such that $\tilde{U}(\tilde{\eta} : [t_1, t_2]) = I$.

Proof. Let I be a weak bi-ideal of M . Let $\tilde{\eta}$ be an i.v fuzzy subset of M defined by

$$\tilde{\eta}(x) = \begin{cases} [t_1, t_2] & \text{if } x \in I \\ \tilde{0} & \text{otherwise} \end{cases}$$

Then $\tilde{U}(\tilde{\eta} : [t_1, t_2]) = I$. Assume that $\tilde{\eta}(x - y) < \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$. This implies that $\tilde{\eta}(x - y) = \tilde{0}$ and $\min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\} = [t_1, t_2]$ so $x, y \in I$ and $\alpha, \beta \in \Gamma$ but $x - y \notin I$, which is a contradiction. Thus, $\tilde{\eta}(x - y) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$. Suppose that $\tilde{\eta}(x\alpha y\beta z) < \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\}$. Then $\tilde{\eta}(x\alpha y\beta z) = \tilde{0}$ and $\min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\} = [t_1, t_2]$ so $x, y, z \in I$ but $x\alpha y\beta z \notin I$ which is a contradiction. Hence $\tilde{\eta}(x\alpha y\beta z) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\}$. ■

Theorem 3.13 Let H be a nonempty subset of M and $\tilde{\eta}$ be an i.v fuzzy subset of M defined by

$$\tilde{\eta}(x) = \begin{cases} \tilde{s} & \text{if } x \in H \\ \tilde{t} & \text{otherwise} \end{cases}$$

for some $x \in M$, $\tilde{s}, \tilde{t} \in D[0, 1]$ and $\tilde{s} > \tilde{t}$. Then H is a weak bi-ideal of M if and only if $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of H .

Proof. Assume that H is a weak bi-ideal of M . Let $x, y \in M$. We consider four Cases:

- (1) $x \in H$ and $y \in H$.
- (2) $x \in H$ and $y \notin H$.
- (3) $x \notin H$ and $y \in H$.
- (4) $x \notin H$ and $y \notin H$.

Case (1): If $x \in H$ and $y \in H$. Then $\tilde{\eta}(x) = \tilde{s} = \tilde{\eta}(y)$. Since H is a weak bi-ideal of M , then $x - y \in H$. Thus, $\tilde{\eta}(x - y) = \tilde{s} = \min^i\{\tilde{s}, \tilde{s}\} = \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$.

Case (2): If $x \in H$ and $y \notin H$. Then $\tilde{\eta}(x) = \tilde{s}$ and $\tilde{\eta}(y) = \tilde{t}$. So, $\min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\} = \tilde{t}$. Now, $\tilde{\eta}(x - y) = \tilde{s}$ or \tilde{t} according as $x - y \in H$ or $x - y \notin H$. By assumption, $\tilde{s} > \tilde{t}$, we have $\tilde{\eta}(x - y) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$. Similarly, we prove Case (3).

Case (4): $x, y \notin H$, we have, $\tilde{\eta}(x) = \tilde{t} = \tilde{\eta}(y)$. So, $\min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\} = \tilde{t}$. Next, $\tilde{\eta}(x - y) = \tilde{s}$ or \tilde{t} , according as $x - y \in H$ or $x - y \notin H$. So, $\tilde{\eta}(x - y) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\}$. Now let $x, y, z \in M$ and $\alpha, \beta \in \Gamma$. We have the following eight Cases.

- (1) $x \in H, y \in H$ and $z \in H$.
- (2) $x \notin H, y \in H$ and $z \in H$.
- (3) $x \in H, y \notin H$ and $z \in H$.
- (4) $x \in H, y \in H$ and $z \notin H$.
- (5) $x \notin H, y \notin H$ and $z \in H$.
- (6) $x \in H, y \notin H$ and $z \notin H$.
- (7) $x \notin H, y \in H$ and $z \notin H$.
- (8) $x \notin H, y \notin H$ and $z \notin H$.

These cases can be proved by arguments similar to the fuzzy cases above. Hence, $\tilde{\eta}(x\alpha y\beta z) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\}$. Hence $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M . Conversely, assume that $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M . Let $x, y, z \in H$ and $\alpha, \beta \in \Gamma$ be such that $\tilde{\eta}(x) = \tilde{\eta}(y) = \tilde{\eta}(z) = \tilde{s}$. Since $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M , we

have $\tilde{\eta}(x - y) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\} = \tilde{s}$ and $\tilde{\eta}(x\alpha y\beta z) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\} = \tilde{s}$. So, $x - y, x\alpha y\beta z \in H$. Hence H is a weak bi-ideal of M . ■

Theorem 3.14 A nonempty subset H of M is a weak bi-ideal of M if and only if the characteristic function f_H is an i.v fuzzy weak bi-ideal of M .

Proof. The proof is straightforward. ■

Theorem 3.15 Let $\tilde{\eta}$ be an i.v fuzzy weak bi-ideal of M then the set $M_{\tilde{\eta}} = \{x \in M \mid \tilde{\eta}(x) = \tilde{\eta}(0)\}$ is weak bi-ideal of M .

Proof. Let $\tilde{\eta}$ be i.v fuzzy weak bi-ideal of M . Let $x, y \in M_{\tilde{\eta}}$. Then $\tilde{\eta}(x) = \tilde{\eta}(0), \tilde{\eta}(y) = \tilde{\eta}(0)$ and $\tilde{\eta}(x - y) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y)\} = \min^i\{\tilde{\eta}(0), \tilde{\eta}(0)\} = \tilde{\eta}(0)$. So $\tilde{\eta}(x - y) = \tilde{\eta}(0)$. Thus $x - y \in M_{\tilde{\eta}}$. For every $x, y, z \in M_{\tilde{\eta}}$ and $\alpha, \beta \in \Gamma$ we have $\tilde{\eta}(x\alpha y\beta z) \geq \min^i\{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\} = \min^i\{\tilde{\eta}(0), \tilde{\eta}(0), \tilde{\eta}(0)\} = \tilde{\eta}(0)$. Thus $x\alpha y\beta z \in M_{\tilde{\eta}}$. Hence $M_{\tilde{\eta}}$ is a weak bi-ideal of M . ■

4. Homomorphism of interval valued fuzzy weak bi-ideals of Γ -near-rings

In this section, we characterize i.v fuzzy weak bi-ideals of Γ -near-rings using homomorphism.

Definition 4.1 [9] Let f be a mapping from a set M to a set S . Let $\tilde{\eta}$ and $\tilde{\delta}$ be i.v fuzzy subsets of M and S respectively. Then $f(\tilde{\eta})$, the image of $\tilde{\eta}$ under f is an i.v fuzzy subset of S defined by

$$f(\tilde{\eta})(y) = \begin{cases} \sup_{x \in f^{-1}(y)}^i \tilde{\eta}(x) & \text{if } f^{-1}(y) \neq \emptyset \\ 0 & \text{otherwise} \end{cases}$$

and the pre-image of $\tilde{\eta}$ under f is an i.v fuzzy subset of M defined by $f^{-1}(\tilde{\delta}(x)) = \tilde{\delta}(f(x))$, for all $x \in M$ and $f^{-1}(y) = \{x \in M \mid f(x) = y\}$.

Definition 4.2 [9] Let M and S be Γ -near-rings. A map $\theta : M \rightarrow S$ is called a (Γ -near-ring)homomorphism if $\theta(x + y) = \theta(x) + \theta(y)$ and $\theta(x\alpha y) = \theta(x)\alpha\theta(y)$ for all $x, y \in M$ and $\alpha \in \Gamma$.

Theorem 4.3 Let $f : M \rightarrow S$ be a homomorphism between Γ -near-rings M and S . If $\tilde{\delta}$ is an i.v fuzzy weak bi-ideal of S , then $f^{-1}(\tilde{\delta})$ is an i.v fuzzy weak bi-ideal of M .

Proof. Let $\tilde{\delta}$ is an i.v fuzzy weak bi-ideal of S . Let $x, y, z \in M$ and $\alpha, \beta \in \Gamma$. Then

$$\begin{aligned} f^{-1}(\tilde{\delta})(x - y) &= \tilde{\delta}(f(x - y)) \\ &= \tilde{\delta}(f(x) - f(y)) \\ &\geq \min^i\{\tilde{\delta}(f(x)), \tilde{\delta}(f(y))\} \\ &= \min^i\{f^{-1}(\tilde{\delta}(x)), f^{-1}(\tilde{\delta}(y))\} \end{aligned}$$

and

$$\begin{aligned}
 f^{-1}(\tilde{\delta})(x\alpha y\beta z) &= \tilde{\delta}(f(x\alpha y\beta z)) \\
 &= \tilde{\delta}(f(x)\alpha f(y)\beta f(z)) \\
 &\geq \min^i\{\tilde{\delta}(f(x)), \tilde{\delta}(f(y)), \tilde{\delta}(f(z))\} \\
 &= \min^i\{f^{-1}(\tilde{\delta}(x)), f^{-1}(\tilde{\delta}(y)), f^{-1}(\tilde{\delta}(z))\}.
 \end{aligned}$$

Therefore $f^{-1}(\tilde{\delta})$ is an i.v fuzzy weak bi-ideal of M . ■

We can also state the converse of the Theorem 4.3 by strengthening the condition on f as follows.

Theorem 4.4 Let $f : M \rightarrow S$ be an onto homomorphism of Γ -near-rings M and S . Let $\tilde{\delta}$ be an i.v fuzzy subset of S . If $f^{-1}(\tilde{\delta})$ is an i.v fuzzy weak bi-ideal of M , then $\tilde{\delta}$ is an i.v fuzzy weak bi-ideal of S .

Proof. Let $x, y, z \in S$. Then $f(a) = x, f(b) = y$ and $f(c) = z$ for some $a, b, c \in M$ and $\alpha, \beta \in \Gamma$. It follows that

$$\begin{aligned}
 \tilde{\delta}(x - y) &= \tilde{\delta}(f(a) - f(b)) \\
 &= \tilde{\delta}(f(a - b)) \\
 &= f^{-1}(\tilde{\delta})(a - b) \\
 &\geq \min^i\{f^{-1}(\tilde{\delta})(a), f^{-1}(\tilde{\delta})(b)\} \\
 &= \min^i\{\tilde{\delta}(f(a)), \tilde{\delta}(f(b))\} \\
 &= \min^i\{\tilde{\delta}(x), \tilde{\delta}(y)\}.
 \end{aligned}$$

and

$$\begin{aligned}
 \tilde{\delta}(x\alpha y\beta z) &= \tilde{\delta}(f(a)\alpha f(b)\beta f(c)) \\
 &= \tilde{\delta}(f(a\alpha b\beta c)) \\
 &= f^{-1}(\tilde{\delta})(a\alpha b\beta c) \\
 &\geq \min^i\{f^{-1}(\tilde{\delta})(a), f^{-1}(\tilde{\delta})(b), f^{-1}(\tilde{\delta})(c)\} \\
 &= \min^i\{\tilde{\delta}(f(a)), \tilde{\delta}(f(b)), \tilde{\delta}(f(c))\} \\
 &= \min^i\{\tilde{\delta}(x), \tilde{\delta}(y), \tilde{\delta}(z)\}.
 \end{aligned}$$

Hence $\tilde{\delta}$ is an i.v fuzzy weak bi-ideal of S . ■

Theorem 4.5 Let $f : M \rightarrow S$ be an onto Γ -near-ring homomorphism. If $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M , then $f(\tilde{\eta})$ is an i.v fuzzy weak bi-ideal of S .

Proof. Let $\tilde{\eta}$ be an i.v fuzzy weak bi-ideal of M . Since $f(\tilde{\eta})(x') = \sup_{f(x)=x'}^i(\tilde{\eta}(x))$ for $x' \in S$ and hence $f(\tilde{\eta})$ is nonempty. Let $x', y' \in S$ and $\alpha, \beta \in \Gamma$. Then we have $\{x|x \in f^{-1}(x' - y')\} \supseteq \{x - y|x \in f^{-1}(x') \text{ and } y \in f^{-1}(y')\}$ and $\{x|x \in f^{-1}(x'y')\} \supseteq$

$$\{x\alpha y | x \in f^{-1}(x') \text{ and } y \in f^{-1}(y')\}.$$

$$\begin{aligned} f(\tilde{\eta})(x' - y') &= \sup_{f(z)=x'-y'}^i \{\tilde{\eta}(z)\} \\ &\geq \sup_{f(x)=x', f(y)=y'}^i \{\tilde{\eta}(x - y)\} \\ &\geq \sup_{f(x)=x', f(y)=y'}^i \{\min^i \{\tilde{\eta}(x), \tilde{\eta}(y)\}\} \\ &= \min^i \{\sup_{f(x)=x'}^i \{\tilde{\eta}(x)\}, \sup_{f(y)=y'}^i \{\tilde{\eta}(y)\}\} \\ &= \min^i \{f(\tilde{\eta})(x'), f(\tilde{\eta})(y')\}. \end{aligned}$$

Next,

$$\begin{aligned} f(\tilde{\eta})(x'\alpha y'\beta z') &= \sup_{f(h)=x'\alpha y'\beta z'}^i \{\tilde{\eta}(h)\} \\ &\geq \sup_{f(x)=x', f(y)=y', f(z)=z'}^i \{\tilde{\eta}(x\alpha y\beta z)\} \\ &\geq \sup_{f(x)=x', f(y)=y', f(z)=z'}^i \{\min^i \{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\}\} \\ &= \min^i \{\sup_{f(x)=x'}^i \{\tilde{\eta}(x)\}, \sup_{f(y)=y'}^i \{\tilde{\eta}(y)\}, \sup_{f(z)=z'}^i \{\tilde{\eta}(z)\}\} \\ &= \min^i \{f(\tilde{\eta})(x'), f(\tilde{\eta})(y'), f(\tilde{\eta})(z')\}. \end{aligned}$$

Therefore $f(\tilde{\eta})$ is an i.v fuzzy weak bi-ideal of S . ■

5. Anti-homomorphism of interval valued fuzzy weak bi-ideals of Γ -near-rings

In this section, we characterize i.v fuzzy weak bi-ideals of Γ -near-rings using anti-homomorphism.

Definition 5.1 [11] Let M and S be Γ -near-rings. A map $\theta : M \rightarrow S$ is called a (Γ -near-ring)anti-homomorphism if $\theta(x + y) = \theta(y) + \theta(x)$ and $\theta(x\alpha y) = \theta(y)\alpha\theta(x)$ for all $x, y \in M$ and $\alpha \in \Gamma$.

Theorem 5.2 Let $f : M \rightarrow S$ be a anti-homomorphism between Γ -near-rings M and S . If $\tilde{\delta}$ is an i.v fuzzy weak bi-ideal of S , then $f^{-1}(\tilde{\delta})$ is an i.v fuzzy weak bi-ideal of M .

Proof. Let $\tilde{\delta}$ be an i.v fuzzy weak bi-ideal of S . Let $x, y, z \in M$ and $\alpha, \beta \in \Gamma$. Then

$$\begin{aligned} f^{-1}(\tilde{\delta})(x - y) &= \tilde{\delta}(f(x - y)) \\ &= \tilde{\delta}(f(y) - f(x)) \\ &\geq \min^i \{\tilde{\delta}(f(y)), \tilde{\delta}(f(x))\} \\ &= \min^i \{\tilde{\delta}(f(x)), \tilde{\delta}(f(y))\} \\ &= \min^i \{f^{-1}(\tilde{\delta}(x)), f^{-1}(\tilde{\delta}(y))\}. \end{aligned}$$

and

$$\begin{aligned}
 f^{-1}(\tilde{\delta})(x\alpha y\beta z) &= \tilde{\delta}(f(x\alpha y\beta z)) \\
 &= \tilde{\delta}(f(z)\alpha f(y)\beta f(x)) \\
 &\geq \min^i\{\tilde{\delta}(f(z)), \tilde{\delta}(f(y)), \tilde{\delta}(f(x))\} \\
 &= \min^i\{\tilde{\delta}(f(x)), \tilde{\delta}(f(y)), \tilde{\delta}(f(z))\} \\
 &= \min^i\{f^{-1}(\tilde{\delta}(x)), f^{-1}(\tilde{\delta}(y)), f^{-1}(\tilde{\delta}(z))\}.
 \end{aligned}$$

Therefore $f^{-1}(\tilde{\delta})$ is an i.v fuzzy weak bi-ideal of M . ■

We can also state the converse of the Theorem 5.2 by strengthening the condition on f as follows.

Theorem 5.3 Let $f : M \rightarrow S$ be an onto anti-homomorphism of Γ -near-rings M and S . Let $\tilde{\delta}$ be an i.v fuzzy subset of S . If $f^{-1}(\tilde{\delta})$ is an i.v fuzzy weak bi-ideal of M , then $\tilde{\delta}$ is an i.v fuzzy weak bi-ideal of S .

Proof. Let $x, y, z \in S$. Then $f(a) = x, f(b) = y$ and $f(c) = z$ for some $a, b, c \in M$ and $\alpha, \beta \in \Gamma$. It follows that

$$\begin{aligned}
 \tilde{\delta}(x - y) &= \tilde{\delta}(f(a) - f(b)) \\
 &= \tilde{\delta}(f(b - a)) \\
 &= f^{-1}(\tilde{\delta})(b - a) \\
 &\geq \min^i\{f^{-1}(\tilde{\delta})(b), f^{-1}(\tilde{\delta})(a)\} \\
 &= \min^i\{\tilde{\delta}(f(a)), \tilde{\delta}(f(b))\} \\
 &= \min^i\{\tilde{\delta}(x), \tilde{\delta}(y)\}
 \end{aligned}$$

and

$$\begin{aligned}
 \tilde{\delta}(x\alpha y\beta z) &= \tilde{\delta}(f(a)\alpha f(b)\beta f(c)) \\
 &= \tilde{\delta}(f(c\alpha b\beta a)) \\
 &= f^{-1}(\tilde{\delta})(c\alpha b\beta a) \\
 &\geq \min^i\{f^{-1}(\tilde{\delta})(c), f^{-1}(\tilde{\delta})(b), f^{-1}(\tilde{\delta})(a)\} \\
 &= \min^i\{f^{-1}(\tilde{\delta})(a), f^{-1}(\tilde{\delta})(b), f^{-1}(\tilde{\delta})(c)\} \\
 &= \min^i\{\tilde{\delta}(f(a)), \tilde{\delta}(f(b)), \tilde{\delta}(f(c))\} \\
 &= \min^i\{\tilde{\delta}(x), \tilde{\delta}(y), \tilde{\delta}(z)\}.
 \end{aligned}$$

Hence $\tilde{\delta}$ is an i.v fuzzy weak bi-ideal of S . ■

Theorem 5.4 Let $f : M \rightarrow S$ be an onto Γ -near-ring anti-homomorphism. If $\tilde{\eta}$ is an i.v fuzzy weak bi-ideal of M , then $f(\tilde{\eta})$ is an i.v fuzzy weak bi-ideal of S .

Proof. Let $\tilde{\eta}$ be an i.v fuzzy weak bi-ideal of M . Since

$f(\tilde{\eta})(x') = \sup_{f(x)=x'}^i \{\tilde{\eta}(x)\}$, for $x' \in S$ and hence $f(\tilde{\eta})$ is nonempty. Let $x', y' \in S$ and $\alpha, \beta \in \Gamma$. Then we have $\{x|x \in f^{-1}(x' - y')\} \supseteq \{x - y|x \in f^{-1}(x') \text{ and } y \in f^{-1}(y')\}$ and $\{x|x \in f^{-1}(x'y')\} \supseteq \{x\alpha y|x \in f^{-1}(x') \text{ and } y \in f^{-1}(y')\}$

$$\begin{aligned} f(\tilde{\eta})(x' - y') &= \sup_{f(z)=x'-y'}^i \{\tilde{\eta}(z)\} \\ &\geq \sup_{f(x)=x', f(y)=y'}^i \{\tilde{\eta}(x - y)\} \\ &\geq \sup_{f(x)=x', f(y)=y'}^i \{\min^i \{\tilde{\eta}(x), \tilde{\eta}(y)\}\} \\ &= \min^i \{\sup_{f(x)=x'}^i \{\tilde{\eta}(x)\}, \sup_{f(y)=y'}^i \{\tilde{\eta}(y)\}\} \\ &= \min^i \{f(\tilde{\eta})(x'), f(\tilde{\eta})(y')\}. \end{aligned}$$

Next,

$$\begin{aligned} f(\tilde{\eta})(x' \alpha y' \beta z') &= \sup_{f(h)=x' \alpha y' \beta z'}^i \{\tilde{\eta}(h)\} \\ &\geq \sup_{f(x)=x', f(y)=y', f(z)=z'}^i \{\tilde{\eta}(x \alpha y \beta z)\} \\ &\geq \sup_{f(x)=x', f(y)=y', f(z)=z'}^i \{\min^i \{\tilde{\eta}(x), \tilde{\eta}(y), \tilde{\eta}(z)\}\} \\ &= \min^i \{\sup_{f(x)=x'}^i \{\tilde{\eta}(x)\}, \sup_{f(y)=y'}^i \{\tilde{\eta}(y)\}, \sup_{f(z)=z'}^i \{\tilde{\eta}(z)\}\} \\ &= \min^i \{f(\tilde{\eta})(x'), f(\tilde{\eta})(y'), f(\tilde{\eta})(z')\}. \end{aligned}$$

Therefore $f(\tilde{\eta})$ is an i.v fuzzy weak bi-ideal of S . ■

Acknowledgements

The second author was supported in part by UGC-BSR Grant #F.25-1/2014-15(BSR)/7-254/2009(BSR) dated 20-01-2015 in India. The third author was supported in part by UGC-BSR Grant # F4-1/2006(BSR)/7-254/2009(BSR) in India.

References

- [1] G. L. Booth, A note on Γ -near-rings, Stud. Sci. Math. Hung. 23 (1988), 471-475.
- [2] Ch. Rao, V. Swaminathan, Anti-homomorphism in fuzzy ideals, International scholarly and scientific research and innovation. 4 (8) (2010), 1211-1214.
- [3] V. Chinnadurai, S. Kadalarasi, Fuzzy weak bi-ideals of near-rings, Annals of Fuzzy Mathematics and Informatics. 11 (2016), 799-808.
- [4] V. Chinnadurai, K. Arulmozhi, S. Kadalarasi, Interval valued fuzzy ideals of Γ -near-rings, Submitted.
- [5] V. Chinnadurai, K. Arulmozhi, S. Kadalarasi, Characterization of fuzzy weak bi-ideals of Γ -near-rings, Submitted.
- [6] V. Chinnadurai, K. Arulmozhi, S. Kadalarasi, Interval valued fuzzy bi-ideals of Γ -near-rings, Submitted.
- [7] V. Chinnadurai, S. Kadalarasi, interval valued fuzzy quasi-ideals of near-rings, Annals of Fuzzy Mathematics and Informatics. 11 (2016), 621-631.
- [8] Y. B. Jun, M. Sapanic, M. A. Ozturk, Fuzzy ideals in gamma near-rings, Tr. J. of Mathematics. 22 (1998), 449-549.
- [9] S. D. Kim, H. S. kim, On fuzzy ideals of near-rings, Bull. Korean. Math. Soc. 33 (1996), 593-601.
- [10] T. Manikatan, Fuzzy bi-ideals of near-rings, Journal of Fuzzy Mathematics. 17 (3) (2009), 659-671.
- [11] M. Murali, k. Rao, B. Venkateswarlu, Anti-fuzzy k-ideals and anti-homomorphism of Γ -semirings, J. International Math.l Virtual Institute. 5 (2015), 37-54.
- [12] N. Meenakumari, T. Tamizh Chelvam, Fuzzy bi-ideals in gamma near-rings, Journal of Algebra Discrete Structures. 9 (1-2) (2011), 43-52.
- [13] Al. Narayanan, T. Manikatan, $(\in, \in \vee q)$ -fuzzy subnear-rings and $(\in, \in \vee q)$ -fuzzy ideals of near-rings, J. Applied Math. Computing. 27 (2004), 983-998.

- [14] G. Pilz, Near-rings, The theory and its applications, North-Holland Publishing Company, Amesterdam, 1983.
- [15] Bh. Satyanarayana, Contributions to near-rings theory Doctoral Thesis, Nagarjuna University, 1984.
- [16] T. Tamizh Chelvam, N. Meenakumari, Bi-ideals of gamma near-rings, Southeast Bulletin of Mathematics. 27 (2004) 983-998.
- [17] N. Thillaigovindan, V. Chinnadurai, Interval valued fuzzy quasi-ideals of semigroups, East Asian Mathematics Journal. 25 (4) (2009), 449-467.
- [18] N. Thillaigovindan, V. Chinnadurai, S. Kadalarasi , Interval valued fuzzy ideals of nera-rings, The Journal of Fuzzy Mathematics. 23 (2) (2015), 471-484.
- [19] N. Thillaigovindan, V. Chinnadurai, S. Coumaressane, T -fuzzy subsemigroups and T -fuzzy ideals of regular Γ -semigroups, Annals of Fuzzy Mathematics and Informatics. 11 (4) (2016), 669-680.
- [20] Y. K. Cho, T. Tamizh Chelvam, S. Jayalakshmi, Weak bi-ideals of near-rings, J. Korean. Soc. Math. Edu. Ser B. Pure Appl. Math. 14 (3) (2007), 153-159.
- [21] L. A. Zadeh, Fuzzy sets, Inform and Control. 8 (1965), 338-353.
- [22] L. A. Zadeh, The concept of a linguistic variable and its application to approximate reasoning, Inform. Sci. 8 (1975), 199-249.