

On characterizations of weakly e -irresolute functions

M. Özkoç^{a,*}, T. Yılmaz^a

^a*Department of Mathematics, Faculty of Science, Muğla Sıtkı Koçman University, 48000
Menteşe-Muğla, Turkey.*

Received 19 February 2017; Revised 28 December 2017; Accepted 28 December 2017.

Communicated by Mohammad Sadegh Asgari

Abstract. The aim of this paper is to introduce and obtain some characterizations of weakly e -irresolute functions by means of e -open sets defined by Ekici [6]. Also, we look into further properties relationships between weak e -irresoluteness and separation axioms and completely e -closed graphs.

© 2017 IAUCTB. All rights reserved.

Keywords: e -irresolute, weakly e -irresolute, strongly e -regular space, completely e -closed graph.

2010 AMS Subject Classification: 54C08, 54C10, 54C05.

1. Introduction

In 1972, Crossley et al. [4] introduced the concept of irresolute functions in topological spaces. The class of α -irresolute functions were introduced by Maheshwari and Thakur [9]. Recently, the class of semi α -irresolute functions and almost α -irresolute functions and weakly B -irresolute functions were introduced in [3], [2] and [14], respectively. In this paper, we introduce and investigate the concept of weakly e -irresolute functions and study several characterizations and some fundamental properties of these classes of functions. Relations between this class and some other existing classes of functions ([5, 6, 10, 12, 13]) are also obtained.

Throughout this paper (X, τ) and (Y, σ) (or simply X and Y) represent nonempty topological spaces on which no separation axioms are assumed unless otherwise stated.

*Corresponding author.

E-mail address: murad.ozkoc@mu.edu.tr (M. Özkoç); tugceyilmaz090@gmail.com (T. Yılmaz).

Let X be a topological space and A be a subset of X . The closure of A and the interior of A are denoted by $cl(A)$ and $int(A)$, respectively. $\mathcal{U}(x)$ denotes all open neighborhoods of the point $x \in X$. A subset A of a space X is called regular open [15] (resp. regular closed [15]) if $A = int(cl(A))$ (resp. $A = cl(int(A))$). The δ -interior [16] of a subset A of X is the union of all regular open sets of X contained in A and is denoted by $int_\delta(A)$. The subset A is called δ -open [16] if $A = int_\delta(A)$, i.e., a set is δ -open if it is the union of regular open sets. The complement of a δ -open set is called δ -closed [16].

The family of all δ -open (resp. δ -closed) sets in X is denoted by $\delta O(X)$ (resp. $\delta C(X)$). A subset A of a space X is called e -open [6] (resp. β -open [1]) if $A \subseteq int(cl_\delta(A)) \cup cl(int_\delta(A))$ (resp. $A \subseteq cl(int(cl(A)))$). The complement of an e -open (resp. β -open) set is said to be e -closed [6] (resp. β -closed [1]). The e -interior [6] of a subset A of X is the union of all e -open sets of X contained in A and is denoted by $e-int(A)$. The e -closure [6] of a subset A of X is the intersection of all e -closed sets of X containing A and is denoted by $e-cl(A)$. The family of all e -open (resp. e -closed, both e -open and e -closed) sets of X is denoted by $eO(X)$ (resp. $eC(X), eR(X)$). The family of all e -open (resp. e -closed, both e -open and e -closed) sets of X containing a point $x \in X$ is denoted by $eO(X, x)$ (resp. $eC(X, x), eR(X, x)$).

We shall use the well-known accepted language almost in the whole of the proofs of theorems in article.

2. Preliminaries

Definition 2.1 [11] A point x of X is called an e - θ -cluster points of $A \subseteq X$ if $e-cl(U) \cap A \neq \emptyset$ for every $U \in eO(X, x)$. The set of all e - θ -cluster points of A is called the e - θ -closure of A and is denoted by $e-cl_\theta(A)$. A subset A is said to be e - θ -closed if and only if $A = e-cl_\theta(A)$. The complement of an e - θ -closed set is said to be e - θ -open. The family of all e - θ -open (resp. e - θ -closed) sets in X is denoted by $e\theta O(X)$ (resp. $e\theta C(X)$).

Theorem 2.2 [6] Let X be a topological space and $A \subseteq X$. Then the followings hold:

- (a) If $A \in eC(X)$, then $A = e-cl(A)$,
- (b) If $A \subseteq B$, then $e-cl(A) \subseteq e-cl(B)$,
- (c) $e-cl(A) \in eC(X)$,
- (d) $x \in e-cl(A)$ if and only if $U \cap A \neq \emptyset$ for each $U \in eO(X, x)$,
- (e) $e-cl(X \setminus A) = X \setminus e-int(A)$.

Theorem 2.3 [11] Let X be a topological space and $A \subseteq X$. Then the followings hold:

- (a) $A \in eO(X)$ if and only if $e-cl(A) \in eR(X)$,
- (b) $A \in eC(X)$ if and only if $e-int(A) \in eR(X)$,
- (c) If $A \in eO(X)$, then $e-cl(A) = e-cl_\theta(A)$,
- (d) $A \in eR(X)$ if and only if $e\theta O(X) \cap e\theta C(X)$,
- (e) $x \in e-cl_\theta(A)$ if and only if $e-cl(U) \cap A \neq \emptyset$ for each $U \in eO(X, x)$,
- (f) $e-int_\theta(X \setminus A) = X \setminus e-cl_\theta(A)$.

Definition 2.4 A function $f : X \rightarrow Y$ is called:

- (a) weakly continuous [8] (briefly *w.c.*) if for each $x \in X$ and for each open set V of Y containing $f(x)$, there exists an open set U of X containing x such that $f[U] \subseteq cl(V)$,
- (b) weakly e -continuous [12] if for each $x \in X$ and for each open set V of Y containing $f(x)$, there exists an e -open set U of X containing x such that $f[U] \subseteq cl(V)$,
- (c) weakly β -continuous [13] if for each $x \in X$ and for each open set V of Y containing


- $f(x)$, there exists a β -open set U of X containing x such that $f[U] \subseteq cl(V)$,
- (d) e -continuous [6] if $f^{-1}[V] \in eO(X)$ for every open set V of Y ,
 - (e) e -irresolute [7] if $f^{-1}[V] \in eO(X)$ for every e -open set V of Y ,
 - (f) β -irresolute [10] if $f^{-1}[V] \in \beta O(X)$ for every β -open set V of Y ,
 - (g) weakly B -irresolute [14] if for each $x \in X$ and for each b -open V of Y containing $f(x)$, there exists a b -open set U of X containing x such that $f[U] \subseteq bcl(V)$.

3. Weakly e -irresolute Functions

In this section we define the notion of weakly e -irresolute functions. Then we obtain several characterizations of them.

Definition 3.1 Let X and Y be topological spaces. A function $f : X \rightarrow Y$ is said to be weakly e -irresolute if for each x in X and for each e -open set V of Y containing $f(x)$, there exists $U \in eO(X, x)$ such that $f[U] \subseteq e-cl(V)$.

Remark 1 We have the following diagram from Definition 2.4 and Definition 3.1. The converses of these implications are not true in general as shown by the following examples.



Example 3.2 Let $X := \{a, b, c\}$, $\tau := \{\emptyset, X, \{a\}, \{c\}, \{a, c\}\}$ and $\sigma := \{\emptyset, X, \{c\}\}$. Define a function $f : (X, \tau) \rightarrow (X, \sigma)$ such that $f(x) = x$. Then f is weakly e -continuous but not weakly e -irresolute.

Example 3.3 Let $X := \{a, b, c, d, e\}$, $\tau := \{\emptyset, X, \{a\}, \{c\}, \{a, c\}, \{c, d\}, \{a, c, d\}\}$. Define a function $f : (X, \tau) \rightarrow (X, \tau)$ such that $f = \{(a, a), (b, d), (c, d), (d, d), (e, e)\}$. Then f is weakly e -irresolute but not e -irresolute.

Remark 2 A weakly e -irresolute function need not be a weakly B -irresolute function as shown by the following example.

Example 3.4 Let $X := \{a, b, c\}$, $\tau := \{\emptyset, X, \{a, b\}\}$. Then $eR(X) = \mathcal{P}(X)$ and $BR(X) = \{\emptyset, \{a\}, \{b\}, \{a, c\}, \{b, c\}, X\}$. Define a function $f : (X, \tau) \rightarrow (X, \tau)$ such that $f = \{(a, b), (b, c), (c, a)\}$. Then f is weakly e -irresolute but not weakly B -irresolute.

QUESTION. Is there any weakly B -irresolute function which is not weakly e -irresolute?

Theorem 3.5 Let $f : X \rightarrow Y$ be a function. Then the following properties are equivalent:

- (a) f is weakly e -irresolute;
- (b) $f^{-1}[V] \subseteq e-int(f^{-1}[e-cl(V)])$ for every $V \in eO(Y)$;
- (c) $e-cl(f^{-1}[V]) \subseteq f^{-1}[e-cl(V)]$ for every $V \in eO(Y)$.

Proof. (a) \implies (b) : Let $V \in eO(Y)$ and $x \in f^{-1}[V]$.

$$\left. \begin{array}{l}
 (V \in eO(Y))(x \in f^{-1}[V]) \implies V \in eO(Y, f(x)) \\
 \text{(a)} \end{array} \right\} \implies$$

$$\implies (\exists U \in eO(X, x))(f[U] \subseteq e-cl(V))$$

$$\begin{aligned}
&\Rightarrow (\exists U \in eO(X, x)) (U \subseteq f^{-1}[e-cl(V)]) \\
&\Rightarrow (\exists U \in eO(X, x)) (x \in U = e-int(U) \subseteq e-int(f^{-1}[e-cl(V)])) \\
&\Rightarrow x \in e-int(f^{-1}[e-cl(V)]). \\
(b) \implies (c) : & \text{ Let } V \in eO(Y) \text{ and } x \notin f^{-1}[e-cl(V)]. \\
& x \notin f^{-1}[e-cl(V)] \Rightarrow f(x) \notin e-cl(V) \\
& \Rightarrow (\exists F \in eO(Y, f(x))) (F \cap V = \emptyset) \\
& \Rightarrow (\exists F \in eO(Y, f(x))) (F \subseteq Y \setminus V) \\
& \Rightarrow (\exists F \in eO(Y, f(x))) (e-cl(F) \subseteq e-cl(Y \setminus V) = Y \setminus V) \\
& \Rightarrow (\exists F \in eO(Y, f(x))) (e-cl(F) \cap V = \emptyset) \\
& \Rightarrow (\exists F \in eO(Y, f(x))) (f^{-1}[e-cl(F) \cap V] = \emptyset) \\
& \Rightarrow (\exists F \in eO(Y, f(x))) (f^{-1}[e-cl(F)] \cap f^{-1}[V] = \emptyset) \\
& \Rightarrow (\exists F \in eO(Y, f(x))) (e-int(f^{-1}[e-cl(F)]) \cap f^{-1}[V] = \emptyset) \\
& \stackrel{(b)}{\Rightarrow} (e-int(f^{-1}[e-cl(F)]) \in eO(X, x)) (e-int(f^{-1}[e-cl(F)]) \cap f^{-1}[V] = \emptyset) \\
& \Rightarrow x \notin e-cl(f^{-1}[V]). \\
(c) \implies (a) : & \text{ Let } x \in X \text{ and } V \in eO(Y, f(x)). \\
& V \in eO(Y, f(x)) \Rightarrow e-cl(V) \in eR(Y, f(x)) \Rightarrow x \notin f^{-1}[e-cl(Y \setminus e-cl(V))] \Big\} \Rightarrow \\
& \hspace{15em} (c) \\
& \Rightarrow x \notin e-cl(f^{-1}[Y \setminus e-cl(V)]) \\
& \Rightarrow (\exists U \in eO(X, x)) (U \cap f^{-1}[Y \setminus e-cl(V)] = \emptyset) \\
& \Rightarrow (\exists U \in eO(X, x)) (f[U] \cap (Y \setminus e-cl(V)) = \emptyset) \\
& \Rightarrow (\exists U \in eO(X, x)) (f[U] \subseteq e-cl(V)). \quad \blacksquare
\end{aligned}$$

Theorem 3.6 Let $f : X \rightarrow Y$ be a function. Then the following properties are equivalent:

- (a) f is weakly e -irresolute;
- (b) $e-cl(f^{-1}[B]) \subseteq f^{-1}[e-cl_\theta(B)]$ for every subset B of Y ;
- (c) $f[e-cl(A)] \subseteq e-cl_\theta(f[A])$ for every subset A of X ;
- (d) $f^{-1}[F] \in eC(X)$ for every e - θ -closed set F of Y ;
- (e) $f^{-1}[V] \in eO(X)$ for every e - θ -open set V of Y .

Proof. (a) \implies (b) : Let $B \subseteq Y$ and $x \notin f^{-1}[e-cl_\theta(B)]$.

$$x \notin f^{-1}[e-cl_\theta(B)] \Rightarrow f(x) \notin e-cl_\theta(B) \Rightarrow (\exists V \in eO(Y, f(x))) (e-cl(V) \cap B = \emptyset) \dots (1)$$

$$V \in eO(Y, f(x)) \Big\} \Rightarrow (\exists U \in eO(X, x)) (f[U] \subseteq e-cl(V)) \dots (2)$$

$$\begin{aligned}
(1), (2) \Rightarrow & (\exists U \in eO(X, x)) (f[U] \cap B = \emptyset) \\
& \Rightarrow (\exists U \in eO(X, x)) (U \cap f^{-1}[B] = \emptyset) \\
& \Rightarrow x \notin e-cl(f^{-1}[B]).
\end{aligned}$$

(b) \implies (c) : Let $A \subseteq X$.

$$A \subseteq X \Rightarrow f[A] \subseteq Y \Big\} \Rightarrow e-cl(A) \subseteq e-cl(f^{-1}[f[A]]) \subseteq f^{-1}[e-cl_\theta(f[A])]$$

$$\Rightarrow f[e-cl(A)] \subseteq e-cl_\theta(f[A]).$$

(c) \implies (d) : Let $F \in e\theta C(Y)$.

$$F \in e\theta C(Y) \Rightarrow f^{-1}[F] \subseteq X \Big\} \Rightarrow f[e-cl(f^{-1}[F])] \subseteq e-cl_\theta(f[f^{-1}[F]]) \subseteq e-cl_\theta(F) =$$

F

$$\Rightarrow e-cl(f^{-1}[F]) \subseteq f^{-1}[F]$$

$$\Rightarrow f^{-1}[F] \in eC(X).$$

(d) \implies (e) : Clear.

(e) \implies (a) : Let $x \in X$ and $V \in eO(Y, f(x))$.

$$\left. \begin{aligned} V \in eO(Y, f(x)) \Rightarrow e-cl(V) \in e\theta O(Y) \\ (e) \end{aligned} \right\} \Rightarrow \\ \Rightarrow (U := f^{-1}[e-cl(V)] \in eO(X, x)) (f[U] = f[f^{-1}[e-cl(V)]] \subseteq e-cl(V)). \quad \blacksquare$$

Theorem 3.7 Let $f : X \rightarrow Y$ be a function. Then the following properties are equivalent:

- (a) f is weakly e -irresolute;
- (b) For each $x \in X$ and each $V \in eO(Y, f(x))$, there exists $U \in eO(X, x)$ such that $f[e-cl(U)] \subseteq e-cl(V)$;
- (c) $f^{-1}[F] \in eR(X)$ for every $F \in eR(Y)$.

Proof. (a) \implies (b) : Let $x \in X$ and $V \in eO(Y, f(x))$.

$$\left. \begin{aligned} V \in eO(Y, f(x)) \\ \text{Theorem 2.3} \end{aligned} \right\} \Rightarrow \left. \begin{aligned} e-cl(V) \in e\theta O(Y) \cap e\theta C(Y) \\ \text{Theorem 3.6(d)(e)} \end{aligned} \right\} \Rightarrow \\ \Rightarrow (U := f^{-1}[e-cl(V)] \in eO(X) \cap eC(X)) (f[e-cl(U)] \subseteq e-cl(V)).$$

(b) \implies (c) : Let $F \in eR(Y)$ and $x \in f^{-1}[F]$.

$$\left. \begin{aligned} (x \in f^{-1}[F])(F \in eR(Y)) \Rightarrow F \in eR(Y, f(x)) \\ (b) \end{aligned} \right\} \Rightarrow \\ \Rightarrow (\exists U \in eO(X, x)) (f[e-cl(U)] \subseteq e-cl[F] = F) \\ \Rightarrow (\exists U \in eO(X, x)) (U \subseteq e-cl(U) \subseteq f^{-1}[F]) \\ \Rightarrow x \in e-int(f^{-1}[F])$$

Then $f^{-1}[F] \in eO(X) \dots (1)$

$$\left. \begin{aligned} (x \in f^{-1}[Y \setminus F])(F \in eR(Y)) \Rightarrow Y \setminus F \in eR(Y, f(x)) \\ (b) \end{aligned} \right\} \Rightarrow \\ \Rightarrow (\exists U \in eO(X, x)) (f[e-cl(U)] \subseteq e-cl[Y \setminus F] = Y \setminus F) \\ \Rightarrow (\exists U \in eO(X, x)) (U \subseteq e-cl(U) \subseteq f^{-1}[Y \setminus F]) \\ \Rightarrow x \in e-int(f^{-1}[Y \setminus F]) \in eO(X)$$

Then $f^{-1}[Y \setminus F] \in eO(X)$ and so $f^{-1}[F] \in eC(X) \dots (2)$

(1), (2) $\implies f^{-1}[F] \in eR(X)$.

(c) \implies (a) : Let $x \in X$ and $V \in eO(Y, f(x))$.

$$\left. \begin{aligned} V \in eO(Y, f(x)) \Rightarrow e-cl(V) \in eR(Y, f(x)) \\ (c) \end{aligned} \right\} \Rightarrow \\ \Rightarrow (U := f^{-1}[e-cl(V)] \in eR(X, x)) (f[U] \subseteq e-cl(V)). \quad \blacksquare$$

Theorem 3.8 Let $f : X \rightarrow Y$ be a function. Then the following properties are equivalent:

- (a) f is weakly e -irresolute;
- (b) $f^{-1}[V] \subseteq e-int_{\theta}(f^{-1}[e-cl_{\theta}(V)])$ for every $V \in eO(Y)$;
- (c) $e-cl_{\theta}(f^{-1}[V]) \subseteq f^{-1}[e-cl_{\theta}(V)]$ for every $V \in eO(Y)$.

Proof. (a) \implies (b) : Let $V \in eO(Y)$.

$$\left. \begin{aligned} V \in eO(Y) \Rightarrow e-cl_{\theta}(V) \in eR(Y) \\ (a) \end{aligned} \right\} \Rightarrow f^{-1}[e-cl_{\theta}(V)] \in eR(X) \\ \Rightarrow f^{-1}[e-cl_{\theta}(V)] \in e\theta O(X) \Rightarrow e-int_{\theta}(f^{-1}[e-cl_{\theta}(V)]) = f^{-1}[e-cl_{\theta}(V)] \supseteq f^{-1}[V].$$

(b) \implies (c) : Let $V \in eO(Y)$.

$$\left. \begin{aligned} V \in eO(Y) \Rightarrow Y \setminus V \in eC(Y) \Rightarrow e-int_{\theta}(Y \setminus V) \in eR(Y) \\ (b) \end{aligned} \right\} \Rightarrow \\ f^{-1}[e-int_{\theta}(Y \setminus V)] \subseteq e-int_{\theta}(f^{-1}[e-cl_{\theta}(e-int_{\theta}(Y \setminus V))]) = e-int_{\theta}(f^{-1}[e-int_{\theta}(Y \setminus V)]) \\ \Rightarrow X \setminus e-int_{\theta}(f^{-1}[e-int_{\theta}(Y \setminus V)]) \subseteq X \setminus f^{-1}[e-int_{\theta}(Y \setminus V)] \\ \Rightarrow e-cl_{\theta}(X \setminus f^{-1}[e-int_{\theta}(Y \setminus V)]) \subseteq f^{-1}[Y \setminus e-int_{\theta}(Y \setminus V)]$$

$$\begin{aligned}
&\Rightarrow e-cl_\theta(f^{-1}[Y \setminus e-int_\theta(Y \setminus V)]) \subseteq f^{-1}[e-cl_\theta(V)] \\
&\Rightarrow e-cl_\theta(f^{-1}[e-cl_\theta(V)]) \subseteq f^{-1}[e-cl_\theta(V)] \\
&\Rightarrow e-cl_\theta(f^{-1}[V]) \subseteq e-cl_\theta(f^{-1}[e-cl_\theta(V)]) \subseteq f^{-1}[e-cl_\theta(V)]. \\
&(c) \implies (a) : \text{Let } V \in eR(Y). \\
&\left. \begin{aligned} V \in eR(Y) &\Rightarrow V \in eO(Y) \\ &\quad (c) \end{aligned} \right\} \Rightarrow e-cl_\theta(f^{-1}[V]) \subseteq f^{-1}[e-cl_\theta(V)] = f^{-1}[V] \\
&\Rightarrow f^{-1}[V] = e-cl_\theta(f^{-1}[V]) \\
&\Rightarrow f^{-1}[V] \in e\theta C(X) \dots (1) \\
&\left. \begin{aligned} V \in eR(Y) &\Rightarrow Y \setminus V \in eR(Y) \Rightarrow Y \setminus V \in eO(Y) \\ &\quad (c) \end{aligned} \right\} \Rightarrow \\
&\Rightarrow e-cl_\theta(f^{-1}[Y \setminus V]) \subseteq f^{-1}[e-cl_\theta(Y \setminus V)] = f^{-1}[Y \setminus V] \\
&\Rightarrow X \setminus f^{-1}[Y \setminus V] \subseteq X \setminus e-cl_\theta(f^{-1}[Y \setminus V]) \\
&\Rightarrow f^{-1}[V] \subseteq e-int_\theta(f^{-1}[V]) \\
&\Rightarrow f^{-1}[V] = e-int_\theta(f^{-1}[V]) \\
&\Rightarrow f^{-1}[V] \in e\theta O(X) \dots (2) \\
&(1), (2) \Rightarrow f^{-1}[V] \in eR(X). \quad \blacksquare
\end{aligned}$$

Theorem 3.9 Let $f : X \rightarrow Y$ be a function. Then the following properties are equivalent:

- (a) f is weakly e -irresolute;
- (b) $e-cl_\theta(f^{-1}[B]) \subseteq f^{-1}[e-cl_\theta(B)]$ for every subset B of Y ;
- (c) $f[e-cl_\theta(A)] \subseteq e-cl_\theta(f[A])$ for every subset A of X ;
- (d) $f^{-1}[F]$ is e - θ -closed in X for every e - θ -closed set F of Y ;
- (e) $f^{-1}[V]$ is e - θ -open in X for every e - θ -open set V of Y .

Proof. (a) \implies (b) : Let $B \subseteq Y$ and $x \notin f^{-1}[e-cl_\theta(B)]$.

$$\left. \begin{aligned} x \notin f^{-1}[e-cl_\theta(B)] &\Rightarrow f(x) \notin e-cl_\theta(B) \Rightarrow (\exists V \in eO(Y, f(x))) (e-cl(V) \cap B = \emptyset) \\ &\quad f \text{ is w.e.i.} \end{aligned} \right\} \Rightarrow$$

$$\begin{aligned}
&\Rightarrow (\exists U \in eO(X, x)) (f[e-cl(U)] \cap B = \emptyset) \\
&\Rightarrow (\exists U \in eO(X, x)) (e-cl(U) \cap f^{-1}[B] = \emptyset) \\
&\Rightarrow x \notin e-cl_\theta(f^{-1}[B]). \\
&(b) \implies (c) : \text{Let } A \subseteq X. \\
&\left. \begin{aligned} A \subseteq X &\Rightarrow f[A] \subseteq Y \\ &\quad (b) \end{aligned} \right\} \Rightarrow e-cl_\theta(A) \subseteq e-cl_\theta(f^{-1}[f[A]]) \subseteq f^{-1}[e-cl_\theta(f[A])] \\
&\Rightarrow f[e-cl_\theta(A)] \subseteq e-cl_\theta(f[A]). \\
&(c) \implies (d) : \text{Let } F \in e\theta C(Y). \\
&\left. \begin{aligned} F \in e\theta C(Y) &\Rightarrow (e-cl_\theta(F) = F) (f^{-1}[F] \subseteq X) \\ &\quad (c) \end{aligned} \right\} \Rightarrow \\
&\Rightarrow f[e-cl_\theta(f^{-1}[F])] \subseteq e-cl_\theta(f[f^{-1}[F]]) \subseteq e-cl_\theta(F) = F \\
&\Rightarrow e-cl_\theta(f^{-1}[F]) \subseteq f^{-1}[F] \\
&\Rightarrow f^{-1}[F] \in e\theta C(X). \\
&(d) \implies (e) : \text{Let } V \in e\theta O(Y). \\
&\left. \begin{aligned} V \in e\theta O(Y) &\Rightarrow Y \setminus V \in e\theta C(Y) \\ &\quad (d) \end{aligned} \right\} \Rightarrow X \setminus f^{-1}[V] = f^{-1}[Y \setminus V] \in e\theta C(X) \\
&\Rightarrow f^{-1}[V] \in e\theta O(X). \\
&(e) \implies (a) : \text{Let } V \in eR(Y). \\
&\left. \begin{aligned} V \in eR(Y) &\Rightarrow V \in e\theta O(Y) \cap e\theta C(Y) \Rightarrow (V \in e\theta O(Y)) (Y \setminus V \in e\theta C(Y)) \\ &\quad (e) \end{aligned} \right\} \Rightarrow \\
&\Rightarrow (f^{-1}[V] \in e\theta O(X)) (X \setminus f^{-1}[V] = f^{-1}[Y \setminus V] \in e\theta O(X))
\end{aligned}$$

$$\Rightarrow (f^{-1}[V] \in e\theta O(X)) (f^{-1}[V] \in e\theta C(X)) \stackrel{\text{Theorem 2.3(d)}}{\Rightarrow} f^{-1}[V] \in eR(X). \quad \blacksquare$$

4. Some Fundamental Properties

Definition 4.1 A topological space X is said to be strongly e -regular if for each point $x \in X$ and each e -open set U of X containing x , there exists $V \in eO(X, x)$ such that $V \subseteq e-cl(V) \subseteq U$.

Theorem 4.2 Let X and Y be topological spaces and $f : X \rightarrow Y$ be a function. If Y is strongly e -regular and $f : X \rightarrow Y$ is weakly e -irresolute, then the function f is e -irresolute.

Proof. $V \in eO(Y)$ and $x \in f^{-1}[V]$.
 $(V \in eO(Y))(x \in f^{-1}[V]) \Rightarrow V \in eO(Y, f(x)) \left. \vphantom{(V \in eO(Y))} \right\} \Rightarrow$
 $\left. \vphantom{(V \in eO(Y))} \right\} \begin{array}{l} Y \text{ is strongly } e\text{-regular} \\ f \text{ is w.e.i.} \end{array} \Rightarrow$
 $\Rightarrow (\exists F \in eO(Y, f(x))) (F \subseteq e-cl(F) \subseteq V) \left. \vphantom{(\exists F \in eO(Y, f(x)))} \right\} \Rightarrow$
 $\Rightarrow (\exists U \in eO(X, x)) (f[U] \subseteq e-cl(F) \subseteq V)$
 $\Rightarrow (\exists U \in eO(X, x)) (U \subseteq f^{-1}[f[U]] \subseteq f^{-1}[e-cl(F)] \subseteq f^{-1}[V])$
 $\Rightarrow x \in e-int(f^{-1}[V])$
 Then $f^{-1}[V] \in eO(X)$. ■

Definition 4.3 A space X is said to be $e-T_2$ [7] if for each pair of distinct points x and y in X , there exist $A \in eO(X, x)$ and $B \in eO(X, y)$ such that $A \cap B = \emptyset$.

Lemma 4.4 [11] A topological space X is $e-T_2$ if and only if for each pair of distinct points x and y of X , there exist $U \in eO(X, x)$ and $V \in eO(X, y)$ such that $e-cl(U) \cap e-cl(V) = \emptyset$.

Theorem 4.5 Let X and Y be topological spaces and $f : X \rightarrow Y$ be a function. If Y is $e-T_2$ and $f : X \rightarrow Y$ is weakly e -irresolute injection, then X is $e-T_2$.

Proof. Let $x, y \in X$ and $x \neq y$.

$$\left. \begin{array}{l} (x, y \in X) (x \neq y) \\ f \text{ is injective} \end{array} \right\} \Rightarrow \left. \begin{array}{l} f(x) \neq f(y) \\ \text{Lemma 4.4} \end{array} \right\} \Rightarrow$$

$$\Rightarrow (\exists V \in eO(Y, f(x))) (\exists W \in eO(Y, f(y))) (e-cl(V) \cap e-cl(W) = \emptyset) \dots (1)$$

$$\left. \begin{array}{l} (V \in eO(Y, f(x))) (W \in eO(Y, f(y))) \\ f \text{ is w.e.i.} \end{array} \right\} \Rightarrow$$

$$\Rightarrow (\exists G \in eO(X, x)) (\exists H \in eO(X, y)) (f[G] \subseteq e-cl(V)) (f[H] \subseteq e-cl(W)) \dots (2)$$

$$(1), (2) \Rightarrow (\exists G \in eO(X, x)) (\exists H \in eO(X, y)) (f[G] \cap f[H] = \emptyset)$$

$$\Rightarrow (\exists G \in eO(X, x)) (\exists H \in eO(X, y)) (f[G \cap H] = \emptyset)$$

$$\Rightarrow (\exists G \in eO(X, x)) (\exists H \in eO(X, y)) (G \cap H = \emptyset)$$

Then X is $e-T_2$. ■

We recall that for a function $f : X \rightarrow Y$, the subset $\{(x, f(x)) \mid x \in X\}$ of the product space $X \times Y$ is called the graph of f and is denoted by $G(f)$.

Definition 4.6 The graph $G(f)$ of a function $f : X \rightarrow Y$ is said to be completely e -closed (briefly $c.e.c.$) if for each $(x, y) \in (X \times Y) \setminus G(f)$, there exist $U \in eO(X, x)$ and $V \in eO(Y, y)$ such that $(e-cl(U) \times e-cl(V)) \cap G(f) = \emptyset$.

Lemma 4.7 The graph of a function $f : X \rightarrow Y$ is completely e -closed if and only if

for each $(x, y) \in (X \times Y) \setminus G(f)$, there exist $U \in eO(X, x)$ and $V \in eO(Y, y)$ such that $f[e-cl(U)] \cap e-cl(V) = \emptyset$.

Proof. *Necessity.* Let $(x, y) \in (X \times Y) \setminus G(f)$.

$$\left. \begin{array}{l} (x, y) \in (X \times Y) \setminus G(f) \\ G(f) \text{ is c.e.c.} \end{array} \right\} \Rightarrow$$

$$\Rightarrow (\exists U \in eO(X, x))(\exists V \in eO(Y, y))([e-cl(U) \times e-cl(V)] \cap G(f) = \emptyset)$$

$$\Rightarrow (\exists U \in eO(X, x))(\exists V \in eO(Y, y))(f[e-cl(U)] \cap e-cl(V) = \emptyset).$$

Sufficiency. Let $(x, y) \in (X \times Y) \setminus G(f)$.

$$\left. \begin{array}{l} (x, y) \in (X \times Y) \setminus G(f) \\ \text{Hypothesis} \end{array} \right\} \Rightarrow (\exists U \in eO(X, x))(\exists V \in eO(Y, y))(f[e-cl(U)] \cap e-cl(V) = \emptyset)$$

$$\Rightarrow (\exists U \in eO(X, x))(\exists V \in eO(Y, y))([e-cl(U) \times e-cl(V)] \cap G(f) = \emptyset). \quad \blacksquare$$

Theorem 4.8 If Y is $e-T_2$ and $f : X \rightarrow Y$ is weakly e -irresolute, then $G(f)$ is completely e -closed.

Proof. Let $(x, y) \in (X \times Y) \setminus G(f)$.

$$\left. \begin{array}{l} (x, y) \in (X \times Y) \setminus G(f) \Rightarrow (x, y) \notin G(f) \Rightarrow y \neq f(x) \\ Y \text{ is } e-T_2 \end{array} \right\} \xrightarrow{\text{Lemma 4.4}} \Rightarrow$$

$$\Rightarrow (\exists V \in eO(Y, f(x)))(\exists W \in eO(Y, y))(e-cl(V) \cap e-cl(W) = \emptyset) \dots (1)$$

$$\left. \begin{array}{l} V \in eO(Y, f(x)) \\ f \text{ is w.e.i.} \end{array} \right\} \xrightarrow{\text{Theorem 3.7(b)}} \Rightarrow (\exists U \in eO(X, x))(f[e-cl(U)] \subseteq e-cl(V)) \dots (2)$$

$$(1), (2) \Rightarrow (\exists U \in eO(X, x))(\exists W \in eO(Y, y))(f[e-cl(U)] \cap e-cl(W) = \emptyset)$$

$$\Rightarrow (\exists U \in eO(X, x))(\exists W \in eO(Y, y))(e-cl(U) \times e-cl(W)) \cap G(f) = \emptyset$$

Then $G(f)$ is completely e -closed. ■

Theorem 4.9 If a function $f : X \rightarrow Y$ is weakly e -irresolute injection and $G(f)$ is completely e -closed, then X is $e-T_2$.

Proof. Let $x, y \in X$ and $x \neq y$.

$$\left. \begin{array}{l} (x, y \in X)(x \neq y) \\ f \text{ is injective} \end{array} \right\} \Rightarrow f(x) \neq f(y) \Rightarrow (x, f(y)) \notin G(f) \left. \begin{array}{l} \xrightarrow{\text{Lemma 4.7}} \\ G(f) \text{ is c.e.c.} \end{array} \right\} \Rightarrow$$

$$\Rightarrow (\exists U \in eO(X, x))(\exists V \in eO(Y, f(y)))(f[e-cl(U)] \cap e-cl(V) = \emptyset) \dots (1)$$

$$\left. \begin{array}{l} V \in eO(Y, f(y)) \\ f \text{ is w.e.i.} \end{array} \right\} \Rightarrow (\exists H \in eO(X, y))(f[H] \subseteq e-cl(V)) \dots (2)$$

$$(1), (2) \Rightarrow (\exists U \in eO(X, x))(\exists H \in eO(X, y))(f[e-cl(U)] \cap f[H] = \emptyset)$$

$$\Rightarrow (\exists U \in eO(X, x))(\exists H \in eO(X, y))(f[e-cl(U) \cap H] = \emptyset)$$

$$\Rightarrow (\exists U \in eO(X, x))(\exists H \in eO(X, y))(e-cl(U) \cap H = \emptyset)$$

$$\Rightarrow (\exists U \in eO(X, x))(\exists H \in eO(X, y))(U \cap H = \emptyset)$$

This means that X is $e-T_2$. ■

Definition 4.10 A topological space X is said to be e -connected [5] if it cannot be written as the union of two nonempty disjoint e -open sets.

Theorem 4.11 If a function $f : X \rightarrow Y$ is weakly e -irresolute surjection and X is e -connected, then Y is e -connected.

Proof. Suppose that Y is not e -connected. Then

$$\Rightarrow (\exists U, V \in eO(Y) \setminus \{\emptyset\})(U \cap V = \emptyset)(U \cup V = Y) \Rightarrow U, V \in eR(Y) \setminus \{\emptyset\} \left. \begin{array}{l} \xrightarrow{\text{Hypothesis}} \\ \end{array} \right\}$$

$$\Rightarrow (f^{-1}[U], f^{-1}[V] \in eR(X) \setminus \{\emptyset\})(f^{-1}[U \cap V] = f^{-1}[\emptyset]) (f^{-1}[U \cup V] = f^{-1}[Y])$$

$$\Rightarrow (f^{-1}[U], f^{-1}[V] \in eR(X) \setminus \{\emptyset\})(f^{-1}[U] \cap f^{-1}[V] = \emptyset)(f^{-1}[U] \cup f^{-1}[V] = X).$$

This means that X is not e -connected. ■

Acknowledgements

This work is financially supported by Muğla Sıtkı Koçman University, Turkey under BAP grant no: 15/181. This work is dedicated to Professor Dr. Gülhan ASLİM on the occasion of her 70th birthday.

References

- [1] M. E. Abd El-Monsef, S. N. El-Deeb, R. A. Mahmoud, β -open sets and β -continuous mappings, Bull-Fac. Sci. Assiut Univ. 12 (1983), 77-90.
- [2] Y. Beceren, Almost α -irresolute functions, Bull. Cal. Math. Soc. 92 (2000), 213-218.
- [3] Y. Beceren, On semi α -irresolute functions, J. Indian. Acad. Math. 22 (2000), 353-362.
- [4] S. G. Crossley, S. K. Hildebrand, Semi-topological properties, Fund. Math. 74 (1972), 233-254.
- [5] E. Ekici, New forms of contra continuity, Carpathian J. Math. 24 (1) (2008), 37-45.
- [6] E. Ekici, On ϵ -open sets, \mathcal{DP}^* -sets and $\mathcal{DP}\mathcal{E}^*$ -sets and decompositions of continuity, Arab. J. Sci. Eng. 33 (2A) (2008), 269-282.
- [7] E. Ekici, Some generalizations of almost contra-super-continuity, Filomat. 21 (2) (2007), 31-44.
- [8] N. Levine, A decomposition of continuity in topological spaces, Amer. Mat. Monthly. 68 (1961), 44-46.
- [9] S. N. Maheshwari, S. S. Thakur, On α -irresolute mappings, Tamkang J. Math. 11 (1980), 209-214.
- [10] R. A. Mahmoud, M. E. Abd El-Monsef, β -irresolute and β -topological invariants, Proc. Pakistan Acad. Sci. 27 (1990), 285-296.
- [11] M. Özkoç, G. Aslm, On strongly θ - e -continuous functions, Bull. Korean Math. Soc. 47 (5) (2010), 1025-1036.
- [12] M. Özkoç, G. Aslm, On weakly e -continuous functions, Hacet. J. Math. Stat. 40 (6) (2011), 781-791.
- [13] V. Popa, T. Noiri, Weakly β -continuous functions, An. Univ. Timioara Ser. Mat.-Inform. 32 (2) (1994), 83-92.
- [14] N. Rajesh, On weakly B -irresolute functions, Acta Univ. Apulensis. 30 (2012), 153-160.
- [15] M. H. Stone, Applications of the theory of Boolean rings to general topology, Trans. Amer. Math. Soc. 41 (1937), 375-481.
- [16] N. V. Veličko, H -closed topological spaces, Amer. Mat. Soc. Transl. Ser. 78 (2) (1968), 103-118.